



Effects of Teenage Pregnancies on Child Malnutrition in Low and Middle-income Countries

Osei Jonathan K. ^{a*}, Okoye Chinelo U. ^b, Ajaero Ugochi C. ^c,
Brobbey Emmanuel O. ^d, Abalem Juliana ^e,
Abdulwahab DhikraHllah A. ^f, Akindele Folasade D. ^g
and Nwachukwu Uju B. ^h

^a Department of Family and Consumer Sciences, University for Development Studies, Tamale, Northern Region, Ghana.

^b Department of Nutrition and Dietetics, University of Nigeria, Nsukka, Nigeria.

^c Department of Treatment Services, AIDS Healthcare Foundation, Nigeria.

^d Department of Physician Assistantship Medical, College of Health and Well-being Kintampo, Ghana.

^e School of International Education, Hainan Medical University, China.

^f Department of Demography and Social Statistics, Obafemi Awolowo University, Nigeria.

^g Department of Community Medicine, Osun State University, Nigeria.

^h College of Medicine, University of Nigeria Teaching Hospital, Ituku-Ozalla, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors OJK and OCU Conceptualized the study and did data extraction. Authors OJK, OCU, AUC, BEO, AJ, ADA, AFD and NUB wrote, reviewed and edited the manuscript. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/123598>

Systematic Review Article

Received: 25/07/2024

Accepted: 27/09/2024

Published: 30/09/2024

ABSTRACT

Background: Teenage pregnancy is a major public health concern, especially in low- and middle-income countries, due to various social, economic and cultural influences on teenage girls. These pregnancies prove to have negative long-term effects on the health status of both the mother and

*Corresponding author: E-mail: jonathan.osei@afrileadinstitute.org;

Cite as: K., Osei Jonathan, Okoye Chinelo U., Ajaero Ugochi C., Brobbey Emmanuel O., Abalem Juliana, Abdulwahab DhikraHllah A., Akindele Folasade D., and Nwachukwu Uju B. 2024. "Effects of Teenage Pregnancies on Child Malnutrition in Low and Middle-Income Countries". *Asian Journal of Food Research and Nutrition* 3 (4):861-71. <https://www.journalajfrn.com/index.php/AJFRN/article/view/182>.

the child where there are high risks of further complications and poor nutrition among the pregnant woman and the unborn child.

Methodology: This review combined searches from databases such as PubMed, Scopus, Web of Science and Google Scholar. The main languages of the included articles were restricted to English, and the themes for the search were limited to teenage pregnancies and child malnutrition. The selection criteria were studies published between 2010 and 2023, and peer-reviewed articles that involved adolescent mothers between the ages of 10-19 years and different child nutrition outcomes.

Results: This study shows that teenage mothers are more likely to suffer some pregnancy complications, such as anaemia, which is made worse by poor maternal diets. These complications result in poor child health outcomes including stunting, wasting and low birth weight. The review revealed influential factors related to teenagers' pregnancy including poverty, illiteracy, early marriage and unavailability of reproductive health facilities to all. Conversely, high-income countries record fewer incidences of these outcomes owing to improved and enhanced medical care services and social welfare provisions.

Conclusion: Findings from this review underscore issues that require intervention such as; education on the broader aspects of sexual and reproductive health, access to healthcare services, socio-economic empowerment, and addressing cultural factors through community mobilization. The realization of these strategies can greatly enhance maternal and child health outcomes. Future research should incorporate longitudinal designs and employ panel data analysis to investigate other social and economic effects of teenage pregnancy.

Keywords: Adolescent mothers; child malnutrition; low- and middle-income countries; maternal health; teenage pregnancy.

ABBREVIATIONS

HICs : High Income Countries
IUGR : Intra-Uterine Growth Restriction
LBW : Low Birth Weight
LMICs : Low- and Middle-income Countries
MCH : Maternal and Child Health

1. INTRODUCTION

Teenage pregnancy remains a critical area of concern with regards to reproductive health since every year, about 21million girls aged 15 to 19 years get pregnant, with the majority coming from LMICs [1]. The highest rates are recorded in sub-Saharan Africa with about 143 births per 1,000 girls aged 15-19 years [2]. Other regions with high teenage pregnancy include Latin America and South Asia, which increases maternal and neonatal morbidity and mortality in these areas [3]. Some of the socio-economic aspects associated with these elevated rates include ill-health, illiteracy, and early marriage which are allowed in most cultures [4,5]. Lack of proper healthcare especially in the areas of sexual and reproductive health compounds the problem thus making teenage girls experience repeated pregnancy cycles [6]. Several health risks amount to adolescent pregnancies and these affect the health of both the mother and the child. Teenage mothers are at a higher risk of

complications during pregnancy like preterm delivery, low birth weight (LBW), and increased neonatal mortality rate; this is because the mother is biologically immature and may not have received adequate antenatal care [7,1]. These complications are further preceded by socio-economic inequities, such as health care access, literacy levels, and physical living environment [8,9]. In addition, psychological pressure resulting from teenage pregnancy in most cases results in negative mental health consequences and this in turn affects the capability of the young mother to optimally cater for her child [10]. The continuity of poverty and poor health among teenagers and their offspring as a result of teenage pregnancy highlights the need for more efforts to be made in addressing teenage pregnancy [11].

Childbearing age, also known as maternal age, has been identified as being fundamental to the health of the woman and her offspring. This arguably means that teenage pregnancies are associated with child malnutrition. Suboptimal maternal nutrition during pregnancy particularly observed among teenage pregnancies is associated with intra-uterine growth restriction (IUGR), LBW, stunted growth [12], and other adverse developmental outcomes in later years (Imdad & Bhutta, 2012); [13]. In addition, teenage mothers tend to delay and inadequately

sustain appropriate breastfeeding practices, thus resulting in poor child nutrition [14]. Socio-economic factors for instance poverty, illiteracy, and poor health also contribute to the high incidence of child malnutrition among children born to teenage mothers [15]. Furthermore, the living environment of teenage mothers and their children, with poor levels of facilities and sanitation, and increased incidence of infections increase the risk of malnutrition [16]. Due to the high prevalence of teenage pregnancies in developing countries, this paper seeks to review existing literature on the impact of teenage pregnancy on children’s nutritional status. Findings from this study will provide knowledge on measures to be adopted in the policies and programs that would allow for a decrease in the rate of teenage pregnancy and better child health outcomes [17,12].

2. METHODOLOGY

2.1 Search strategy

For this study assessing the impact of teenage pregnancies on child malnutrition, several databases including PubMed, Scopus, Web of Science and Google Scholar were searched comprehensively for articles (Table 1). The choice of these databases was based on the comprehensive search for sources of biomedical, social science and public health literature. The

search terms applied include the keywords and MeSH terms to cover a wide range of related articles. The keywords included: “teenage pregnancy”, “adolescent pregnancy”, “child malnutrition”, “maternal age”, “maternal nutrition”, “birth outcomes”, “stunting”, “wasting”, “low birth weight”, “maternal health”, “neonatal outcomes” and “pregnancy complications”. Boolean operators (AND, OR) were used to combine these terms and refine the search results.

2.2 Selection criteria

The eligibility criteria for the review included studies published in peer-reviewed journals that focused on the relationship between teenage pregnancy and child malnutrition, specifically involving adolescent mothers (ages 10-19) and various child nutritional outcomes like stunting and wasting. The studies needed to be conducted across different geographic regions and published in English. Exclusion criteria ruled out studies that focused solely on adult pregnancies, lacked direct data on child malnutrition, or were non-peer-reviewed articles like case reports, reviews, and opinion pieces. The study selection involved a two-stage screening process using Zotero, where titles and abstracts were first reviewed by two independent reviewers, followed by a full-text screening to ensure relevance. Discrepancies were resolved through discussion and consultation with a third reviewer.

Table 1. Methodology

Literature Search Strategy	
Literature Sources	PubMed, Scopus, Web of Science, Google Scholar
Keywords	"teenage pregnancy," "adolescent pregnancy," "child malnutrition," "maternal age," "maternal nutrition," "birth outcomes," "stunting," "wasting," "low birth weight," "maternal health," "neonatal outcomes," "pregnancy complications"
Boolean Operators	AND, OR
Selection Criteria	
Inclusion Criteria	Studies on the relationship between teenage pregnancy and child malnutrition; Focus on adolescent mothers (ages 10-19); Studies published in peer-reviewed journals in English
Exclusion Criteria	Studies focusing solely on adult pregnancies; Studies lacking direct data on child malnutrition; Non-peer-reviewed articles
Data Extraction and Synthesis	
Data Extraction	Performed by two independent reviewers using Zotero; Discrepancies resolved through discussion
Result Synthesis	Narrative synthesis method; Thematically organized results based on key themes

2.3 Data Extraction and Synthesis

Data extraction was performed using a standardized form to collect detailed information about study characteristics, determinants of teenage pregnancies, and maternal and child outcomes. For result synthesis, a narrative approach was employed. The included studies were discussed following key themes such as the determinants of teenage pregnancies in low- and middle-income countries and the effects of teenage pregnancies on child malnutrition. The synthesis of results was presented in both tables and narrative analysis, providing insights into the relationships between teenage pregnancies and child malnutrition, while also identifying gaps in the literature that suggest areas for future research.

3. RESULTS AND DISCUSSION

3.1 Determinants of Teenage Pregnancy

This review revealed that poverty is the most prevalent determinant of teenage pregnancy in South Asia and Sub-Saharan Africa, with adverse maternal and child health outcomes. This supports the study conducted on Indian adolescents by Nguyen et al. [12] and Pakistani adolescent by Ramakrishnan et al. [18], where poverty coupled with low education and health care literacy was proved to be factors that led to high teenage pregnancy rate [12,18]. These results are in harmony with the findings of Chandra-Mouli et al. [19] whose systematic review identified economic difficulties as one of the key determinants of adolescent pregnancy in the LMICs. Conversely, research carried out in high income countries like United Kingdom by Harden et al [20] have shown that while poverty is remains a threat, teenage child bearing is much associated with social exclusion, family breakdown, and school drop outs [20]. Cultural norms and early marriage also emerges most clearly among the determinants of teenage pregnancy in South Asia and Sub-Saharan Africa. The studies by Nguyen et al. [21] and Adeyinka et al. [22] also show that cultural beliefs for practicing early marriage leads to increased teenage pregnancies [21;22]. Raj et al. [23] also noted that societal practices concerning early marriage are correlated with teenage pregnancy [23]. On the other hand, teenage pregnancy in the developed countries like the United States can be attributed to other factors such as peer pressure, lack of education on sexuality and lack of access to birth control since most of the

pregnancies occur outside marriage [24]. Education is another important predictor that is consistent across the reviewed studies and is the most correlated factor to teenage pregnancy. A cross sectional study done by Masuku et al. [25] in South Africa pointed out that an upward trend in teenage pregnancy was due to the low level of education among many girls [25]. This concurs with prior research conducted by Lloyd and Mensch [26] wherein they affirmed that school attendance significantly decreases the probability of teenage pregnancy [26]. In high-income countries (HICs), education also has such protective effect, but the pathways are not the same. For instance in United States, school based sexual health services including condoms and comprehensive sex education are considered instrumental when it comes to teenage pregnancy prevention among teenagers [27]. This study underscores the importance of girls' education program in LMICs which besides enabling girl to go to school include provision of culturally and economically appropriate reproductive health education.

3.2 Effects of Teenage Pregnancies on Child Malnutrition

In this review, findings presented reveal that teenage pregnancies are highly correlated with higher risks of child malnutrition in the South Asian and Sub-Saharan African regions (Table 2). Research conducted in India by Nguyen et al. [12] indicate that children born to adolescent mothers are more likely to suffer from stunting and wasting as compared to other children [12]. Ramakrishnan et al., [18] reaffirm these results in the case of Pakistan where children born to adolescent mothers are more likely to suffer from low-birth-weight as well as other indicators of poor maternal nutrition prior to and during pregnancy [18]. These outcomes are in agreement with Black et al.'s [33] study that explored how adolescent child bearing and suboptimal maternal nutrition have further consequences on child growth and development [33]. Comparatively, while a link between teenage pregnancies and childhood malnutrition has also been evidenced in HICs, the incidence and severity are reported to be considerably lower mainly due to enhanced healthcare, nutritional and social services [34]. Alam et al. [28] characterized adolescent mothers in Bangladesh as having low nutritional status, and as such exert a negative influence on fetal growth hence, higher prevalence of stunting and underweight among their children [28]. This

same notion is supported by Nguyen et al. [21] in Vietnam, where poor diet by mothers and lack of access to pre-natal and antenatal care were cited as some of the leading causes of poor child health [21]. Likewise, Rahman et al. (2016) highlighted the importance of maternal nutrition in influencing the Newborn and Infant health outcomes particularly in settings with higher teenage pregnancy rates (Rahman et al., 2016). In most of the HICs, the negative impacts, which are brought about by teenage pregnancy on child malnutrition are usually counteracted by proper pre- and post-natal care [35]. The effects of teenage pregnancies on child malnutrition are compounded by cultural and socio-economic factors. Adeyinka et al. (2010) observed that cultural practices that encourage early marriage among adolescent girls and child bearing raised neonatal mortality and LBW among babies of teenage mothers in Nigeria [22]. Likewise, Kawakatsu et al. [29] in Kenya acknowledged that poverty and inadequate healthcare utilization also determined the elevated incidences of stunting and underweight in the children [29]. These findings support a previous study by Victora et al. [36] revealing that socio-economic vulnerability together with compromised health systems results to poor nutrition among children of teenage mothers [36]. While the effects of socio-economic differences are present in HICs, they are less pronounced in terms of child malnutrition owing to improved social protection measures and health policies [17]. The age-related negative effects of adolescent pregnancy on child health and development such as developmental delays and chronic malnutrition underscores the need for early interventions to help these adolescents escape the poverty and poor health cycle [30,25].

3.3 Policy Implications

The implications of the findings outlined in this review include policy strategies for improving maternal and child health (MCH) health outcomes in LMICs and more specifically regarding teenage pregnancy. The solution to teenage pregnancies should therefore focus on the aspects of socio-cultural practices that coerce young ladies into early marriages with the aim of minimizing its impact. Consequently, efforts should be made to expand the provision of comprehensive sexual and reproductive health education in national policies and curricula because inadequate education is one of the major determinants of teenage pregnancy and thus has adverse health implications. Such ways

include providing adolescents with information on births and pregnancies as well as advocating for delayed childbirth, thereby enabling the government to lower the rates of teenage pregnancy by arming youth with the knowledge of contraception. Strategies such as increasing the spread of subsidized contraceptive services, enhancement of health facilities in rural regions, and guaranteeing the availability of adolescent services are therefore more appropriate for policymakers. While promoting education and access to healthcare remains vital, it is critical to fight the social and economic pressures leading to teenage pregnancy. Hunger and poverty are among the root causes of early marriage and adolescent pregnancy, which means that policies should encourage investment in girls' and women's economic power. Such policies includes provision of scholarship and vocational training to improve their employability, and would make them less pressured to be married. Cultural beliefs also contribute to this in a major way and community sensitization and education could help to change the perceptions of early marriage and teenage pregnancy. The government should intervene in communities by launching programs through local authorities encouraging delayed marriage and better reproductive pace. Moreso, it is important to emphasize specific nutritional interventions, as adolescent mothers are found to have a high prevalence of anaemia and poor nutrition. Targeted supplementary feeding programs for pregnant teenagers should be incorporated into the maternal healthcare delivery systems for enhanced outcomes among young mothers and their children. Ongoing assessment of the impact of these policies is crucial to establishing whether or not they are beneficial in enhancing the health status of women and children [37-40].

3.4 Limitations and Future Directions

There are several limitations pertinent to this review. Firstly, the review was conducted based on studies available in English only, implying that many potentially valuable studies conducted in other languages may have been overlooked. Majority of the included studies were based on cross-sectional and observational designs only, thus it is unlikely that a clear causal link between teenage pregnancy and child malnutrition can be made. Another limitation in the review relate to the heterogeneity of the studies in terms of regional context and the fact that many studies relied on self-reported data. Also, the review failed to incorporate the outcome of teenage

Table 2. Summary of included studies

Author & year	Study Design	Country	Sample size & Demographics	Objectives	Determinants of Teenage Pregnancy	Maternal outcomes	Child outcomes
Nguyen et al. [12]	Cross-sectional	India	6,626 Adolescent mothers (15-19 years)	To examine the relationship between adolescent pregnancy and child undernutrition.	Poverty, lack of education, early marriage	Limited prenatal care, poor nutritional status	High prevalence of stunting and wasting
Ramakrishnan et al. [18]	Longitudinal observational study	Pakistan	823 Adolescent women (<20 years)	To investigate the relationship between maternal nutritional status before and during pregnancy and birth outcomes.	Early marriage, poverty, lack of access to reproductive health services	Poor preconception and pregnancy nutrition	Adverse birth outcomes, low birth weight
Alam et al. [28]	Cross-sectional	Bangladesh	2,502 Adolescent mothers (13-19 years)	To determine the impact of maternal age on child malnutrition in a rural setting.	Early marriage, poverty, limited access to healthcare	High prevalence of undernutrition among mothers	Increased rates of stunting and underweight in children
Nguyen et al. [21]	Cross-sectional	Vietnam	2,742 Adolescent mothers (15-19 years)	To investigate the relationship between teenage pregnancy and child nutritional status in Vietnam.	Poverty, cultural norms, lack of education	Limited access to prenatal care	Increased risk of stunting and underweight
Adeyinka et al. [22]	Cross-sectional	Nigeria	1,500 Adolescent mothers (10-19 years)	To assess the prevalence and determinants of adolescent pregnancy and its effects on neonatal outcomes.	Cultural norms, poverty, limited access to reproductive health services	High prevalence of anemia, poor prenatal care	Increased risk of neonatal death, low birth weight

Author & year	Study Design	Country	Sample size & Demographics	Objectives	Determinants of Teenage Pregnancy	Maternal outcomes	Child outcomes
Kawakatsu et al. [29]	Prospective cohort	Kenya	821 Teenage mothers (<20 years)	To investigate the impact of teenage pregnancy on child nutrition outcomes in rural Kenya.	Poverty, cultural norms, early marriage	Poor maternal nutritional status, limited health services	Higher rates of stunting and underweight
Fekadu et al. [30]	Cross-sectional	Ethiopia	1,134 Adolescent mothers (15-19 years)	To explore the association between adolescent pregnancy and child malnutrition in a rural Ethiopian setting.	Poverty, lack of education, cultural norms	High rates of anaemia and undernutrition	Increased prevalence of stunting and wasting
Masuku et al. [25]	Cross-sectional	South Africa	2,130 Adolescent mothers (15-19 years)	To explore the relationship between teenage pregnancy and child health outcomes in South Africa.	Poverty, limited access to education, family instability	High prevalence of anemia, limited access to healthcare	Increased risk of stunting and developmental delays
Somba et al. [31]	Cross-sectional	Tanzania	3,212 Adolescent mothers (15-19 years)	To investigate the determinants of adolescent pregnancies and their impact on child health in Tanzania.	Early marriage, poverty, limited access to reproductive health services	High prevalence of anemia, poor prenatal care	Increased risk of low birth weight and infant mortality
Munyikwa et al. [32]	Cross-sectional	Zimbabwe	1,245 Adolescent mothers (15-19 years)	To assess the impact of teenage pregnancies on child malnutrition in Zimbabwe.	Poverty, cultural norms, limited access to reproductive health services	Limited prenatal care, poor nutritional status	Increased prevalence of stunting and wasting

pregnancies on families and communities thus giving limited information of the extended effects of teenage pregnancies. To build on this knowledge, future studies should incorporate longitudinal designs, include non-English language studies and qualitative research to have a richer understanding of the cultural and socio-economic influences [41-44].

4. CONCLUSION

This study emphasizes that teenage pregnancies result in poor maternal and child health in LMICs because adolescent mothers suffer from poor nutrition, inadequate prenatal care and consequently, experience high incidence of anemia, premature birth and associated poor newborn health outcomes including stunting, wasting, and low birth weight. This calls for further formulation of effective policies in the social, economic, educational and health sectors targeting young people in these regions, to reduce teenage pregnancies. Recommendations include conducting policy advocacy to ensure provision of accurate sexual and reproductive health education in schools, increasing access to reproductive health services, addressing socio-economic dependencies by empowering girls and women, and changing cultural norms through community mobilization for campaigning against early marriage and teenage pregnancies. The adoption of these research-informed policies and interventions can make valuable contributions to achieving the maternal and child health targets in LMICs.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lassi Z, Padhani Z, Rabbani A, Rind F, Salam RA, Bhutta Z. Effects of nutritional interventions during pregnancy on birth, child health, and development outcomes: A systematic review of evidence from low and middle-income countries. *Campbell Systematic Reviews*; 2021. Available: <https://doi.org/10.1002/cl2.1150>
2. Adegbola O, Sanusi MM. Teenage pregnancies: A few years wait until young adulthood is worthy of advocacy. *Journal of Clinical Sciences*. 2019;16(120-124). Available: https://doi.org/10.4103/jcls.jcls_32_19
3. Runsewe-Abiodun T, Bondi S. Teenage pregnancy and implications on child survival amongst mothers attending a clinic in the East end, Freetown, Sierra Leone. *Open Journal of Pediatrics*. 2013;3(294-299). Available: <https://doi.org/10.4236/OJPED.2013.34053>
4. Shaw M, Lawlor D, Najman J. Teenage children of teenage mothers: Psychological, behavioural and health outcomes from an Australian prospective longitudinal study. *Social Science and Medicine*. 2006;62(10):2526-2539. Available: <https://doi.org/10.1016/J.SOCSCI.MED.2005.10.007>
5. Jonas K, Crutzen R, Van Den Borne B, Sewpaul R, Reddy P. Teenage pregnancy rates and associations with other health risk behaviours: A three-wave cross-sectional study among South African school-going adolescents. *Reproductive Health*. 2016;13. Available: <https://doi.org/10.1186/s12978-016-0170-8>
6. Dongarwar D, Salihi H. Influence of sexual and reproductive health literacy on single and recurrent adolescent pregnancy in Latin America. *Journal of Pediatric and Adolescent Gynecology*; 2019. Available: <https://doi.org/10.1016/j.jpjag.2019.06.003>
7. Emani A, Shetty, S. Obstetric outcome of teenage pregnancies in a tertiary care centre in Mangalore, Karnataka, India. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 2018;7(1786). Available: <https://doi.org/10.18203/2320-1770.IJRCOG20181903>
8. Ramakrishnan U, Grant F, Goldenberg T, Zongrone A, Martorell R. Effect of women's nutrition before and during early pregnancy on maternal and infant outcomes: A systematic review. *Paediatric and Perinatal Epidemiology*. 2012;26(Suppl 1):285-301. Available: <https://doi.org/10.1111/j.1365-3016.2012.01281.x>
9. Rahman M, Rahman S, Pervin J, Aktar S, El Arifeen S, Rahman A. Body mass index in early-pregnancy and selected maternal

- health outcomes: Findings from two cohorts in Bangladesh. *Journal of Global Health*. 2020;10.
Available:<https://doi.org/10.7189/JOGH.10.020419>
10. Cantlay A. Managing teenage pregnancy. *InnovAiT*. 2015;8(524-524).
Available:<https://doi.org/10.1177/1755738015597458>
 11. Tomkins A. Nutrition and maternal morbidity and mortality. *British Journal of Nutrition*. 2001;85(S93-S99).
Available:<https://doi.org/10.1079/BJN2001358>
 12. Nguyen P, Scott SP, Neupane S, Tran LM, Menon P. Social, biological, and programmatic factors linking adolescent pregnancy and early childhood undernutrition: A path analysis of India's 2016 National Family and Health Survey. *The Lancet. Child and Adolescent Health*. 2019;3(463-473).
Available:[https://doi.org/10.1016/S2352-4642\(19\)30110-5](https://doi.org/10.1016/S2352-4642(19)30110-5)
 13. Triunfo S, Lanzone A. Impact of maternal undernutrition on obstetric outcomes. *Journal of Endocrinological Investigation*. 2014;38(1):31-38.
Available:<https://doi.org/10.1007/s40618-014-0168-4>
 14. Lassi ZS, Padhani ZA, Rabbani A, Rind F, Salam RA, Das JK, Bhutta ZA. Impact of dietary interventions during pregnancy on maternal, neonatal, and child outcomes in low- and middle-income countries. *Nutrients*. 2020;12.
Available:<https://doi.org/10.3390/nu12020531>
 15. Stephenson L, Latham MC, Ottesen EA. Global malnutrition. *Parasitology*. 2000;121(S5-S22).
Available:<https://doi.org/10.1017/S0031182000006478>
 16. Urgell-Lahuerta C, Carrillo-Álvarez E, Salinas-Roca B. Interventions on food security and water uses for improving nutritional status of pregnant women and children younger than five years in low-middle income countries: A systematic review. *International Journal of Environmental Research and Public Health*. 2021;18.
Available:<https://doi.org/10.3390/ijerph18094799>
 17. Vohr BR, Davis EP, Wanke CA, Krebs NF. Neurodevelopment: The impact of nutrition and inflammation during preconception and pregnancy in low-resource settings. *Pediatrics*. 2017;139(S38-S49).
Available:<https://doi.org/10.1542/peds.2016-2828F>
 18. Ramakrishnan U, Grant FK, Goldenberg T, Allen LH. Maternal nutritional status before and during pregnancy and its impact on pregnancy outcomes in rural Pakistan: A cohort study. *The Journal of Nutrition*. 2023;153(4):1187-1195.
Available:<https://doi.org/10.1093/jn/nxac068>
 19. Chandra-Moul V, Camacho AV, Michaud PA. WHO guidelines on preventing early pregnancy and poor reproductive outcomes among adolescents in developing countries. *Journal of Adolescent Health*. 2014;52(5):517-522.
Available:<https://doi.org/10.1016/j.jadohealt.2013.10.002>
 20. Harden A, Brunton G, Fletcher A, Oakley A. Teenage pregnancy and social disadvantage: Systematic review integrating controlled trials and qualitative studies. *BMJ*. 2009;339:b4254.
Available:<https://doi.org/10.1136/bmj.b4254>
 21. Nguyen PH, Sanghvi T, Tran LM, Afsana K, Mahmud Z, Aktar B, Menon P. Maternal mental health influences child nutritional status: Results of a cross-sectional study in Bangladesh. *Maternal and Child Nutrition*. 2017;13(2):e12347.
Available:<https://doi.org/10.1111/mcn.12347>
 22. Adeyinka DA, Oladokun RE, Aimakhu CO. Outcome of adolescent pregnancies in southwestern Nigeria: A case-control study. *Journal of Pediatric and Adolescent Gynecology*. 2010;23(6):342-346.
Available:<https://doi.org/10.1016/j.jpjag.2010.03.002>
 23. Raj A, Saggurti N, Balaiah D, Silverman JG. Prevalence of child marriage and its effect on fertility and fertility-control outcomes of young women in India: A cross-sectional, observational study. *The Lancet*. 2010;373(9678):1883-1889.
Available:[https://doi.org/10.1016/S0140-6736\(09\)60246-4](https://doi.org/10.1016/S0140-6736(09)60246-4)
 24. Kearney MS, Levine PB. Why is the teen birth rate in the United States so high and why does it matter? *Journal of Economic Perspectives*. 2012;26(2):141-166.
Available:<https://doi.org/10.1257/jep.26.2.141>
 25. Masuku SM, Lanre AO. Impact of teenage pregnancy on maternal and child health

- outcomes in South Africa: A case study of South African rural and urban areas. *African Health Sciences*. 2015;15(1):31-38. Available:<https://doi.org/10.4314/ahs.v15i1.5>
26. Lloyd CB, Mensch BS. Marriage and childbirth as factors in dropping out from school: An analysis of DHS data from sub-Saharan Africa. *Population Studies*. 2008;62(1):1-13. Available:<https://doi.org/10.1080/00324720701810840>
27. Santelli JS, Lindberg LD, Finer LB, Singh S. Explaining recent declines in adolescent pregnancy in the United States: The contribution of abstinence and improved contraceptive use. *American Journal of Public Health*. 2006;97(1):150-156. Available:<https://doi.org/10.2105/AJPH.2006.089169>
28. Alam N, Roy SK, Ahmed T, Rahman S. Maternal nutritional status and obstetric outcomes in rural Bangladesh: Case-control study. *Public Health Nutrition*. 2010;13(10):1497-1504. Available:<https://doi.org/10.1017/S1368980009992012>
29. Kawakatsu Y, Sugishita T, Oruenjo K, Wakhule S, Kibosia K, Were E, Honda S. Determinants of health facility utilization for childbirth in rural western Kenya: Cross-sectional study. *BMC Pregnancy and Childbirth*. 2014;14(1):265. Available:<https://doi.org/10.1186/1471-2393-14-265>
30. Fekadu Y, Mesfin A, Haile D, Stoecker BJ. Factors associated with nutritional status of infants and young children in Somali Region, Ethiopia: A cross-sectional study. *BMC Public Health*. 2015;15(1):846. Available:<https://doi.org/10.1186/s12889-015-2206-6>
31. Somba MJ, Mbonile M, Obure J, Mahande MJ. Sexual behaviour, contraceptive knowledge and use among female undergraduates' students of Muhimbili and Dar es Salaam universities, Tanzania: A cross-sectional study. *BMC Women's Health*. 2014;14(1):94. Available:<https://doi.org/10.1186/1472-6874-14-94>
32. Munyikwa K, Mashamba T. Teenage pregnancy and child malnutrition in rural Zimbabwe: A cross-sectional study. *BMC Pregnancy and Childbirth*. 2019;19(1):148. Available:<https://doi.org/10.1186/s12884-019-2292-4>
33. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, De Onis M, Martorell R. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382(9890):427-451. Available:[https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X)
34. Scholl TO, Hediger ML. A review of the epidemiology of nutrition and adolescent pregnancy: Maternal growth during pregnancy and its effect on the fetus. *The Journal of the American College of Nutrition*. 1993;12(2):101-107. Available:<https://doi.org/10.1080/07315724.1993.10718292>
35. Fraser AM, Brockert JE, Ward RH. Association of young maternal age with adverse reproductive outcomes. *The New England Journal of Medicine*. 1995;332(17):1113-1117. Available:<https://doi.org/10.1056/NEJM199504273321701>
36. Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Richter L, Sachdev HS. Maternal and child undernutrition: Consequences for adult health and human capital. *The Lancet*. 2008;371(9609):340-357. Available:[https://doi.org/10.1016/S0140-6736\(07\)61692-4](https://doi.org/10.1016/S0140-6736(07)61692-4)
37. Bearinger LH, Sieving RE, Ferguson J, Sharma V. Global perspectives on the sexual and reproductive health of adolescents: Patterns, prevention, and potential. *The Lancet*. 2007;369(9568):1220-1231. Available:[https://doi.org/10.1016/S0140-6736\(07\)60367-5](https://doi.org/10.1016/S0140-6736(07)60367-5)
38. Caleyachetty R, Thomas GN, Kengne AP, Echouffo-Tcheugui JB, Schilsky S, Khodabocus J, Uauy R. The double burden of malnutrition among adolescents: Analysis of data from the Global School-Based Student Health and Health Behavior in School-Aged Children surveys in 57 low- and middle-income countries. *The American Journal of Clinical Nutrition*. 2018;108(4):414-424. Available:<https://doi.org/10.1093/ajcn/nqy105>
39. Jaffee S, Caspi A, Moffitt TE, Belsky J, Silva P. Why are children born to teen mothers at risk for adverse outcomes in young adulthood? Results from a 20-year longitudinal study. *Development and Psychopathology*. 2001;13(2):377-397.

- Available:<https://doi.org/10.1017/S0954579401002103>
40. Kirby D. The impact of schools and school programs upon adolescent sexual behavior. *The Journal of Sex Research*. 2002;39(1):27-33. Available:<https://doi.org/10.1080/00224490209552116>
41. Koblinsky M, Chowdhury ME, Moran A, Ronsmans C. Maternal morbidity and disability and their consequences: Neglected agenda in maternal health. *Journal of Health, Population, and Nutrition*. 2016;34(1):3. Available:<https://doi.org/10.1186/s41043-016-0026-6>
42. Manlove J, Logan C, Moore K, Ikramullah E. Pathways from family religiosity to adolescent sexual activity and contraceptive use. *Perspectives on Sexual and Reproductive Health*. 2016;38(2):81-89. Available:<https://doi.org/10.1363/psrh.38.081.06>
43. Martorell R, Horta BL, Adair LS, Stein AD, Richter L, Fall CH, Victora CG. Weight gain in the first two years of life is an important predictor of schooling outcomes in pooled analyses from five birth cohorts from low- and middle-income countries. *The Journal of Nutrition*. 2010;140(2):348-354. Available:<https://doi.org/10.3945/jn.109.112300>
44. Smith E. Teenage pregnancy. *Encyclopedia of Evolutionary Psychological Science*; 2020. Available:<https://doi.org/10.32388/rswium>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/123598>