



Effect of Spacing and Nitrogen Management on Growth and Yield of Summer Sesame (*Sesamum indicum* L.)

**Vaka Madhavi^{a++*}, Rajesh Singh^{a#}, Thakur Indu^{a†},
Akankhya Pradhan^{a†} and Dondapati Suneeth^{a++}**

^a Department of Agronomy, Faculty of Agriculture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i51756

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97160>

Original Research Article

Received: 30/12/2022

Accepted: 01/03/2023

Published: 24/03/2023

ABSTRACT

A field experiment was carried out in *Zaid* 2022 at the Department of Agronomy's Agricultural Research Farm in Prayagraj (U.P). The three planting geometries used in the experiment are 30 cm x 10 cm, 40 cm x 10 cm, and 50 cm x 10 cm. There are three nitrogen management treatments used in the experiment: 50% of the Recommended Dose of Nitrogen (RDN) through inorganic fertiliser, 50% of the RDN through inorganic fertiliser in combination with 50% of Nitrogen provided through FYM with *Azotobacter* seed inoculation, and 50% of the RDN through inorganic fertiliser. Ten treatments were duplicated three times in the randomised block design of the experiment.

⁺⁺ M. Sc. Scholar;

[#] Associate Professor;

[†] PhD Scholar;

*Corresponding author: E-mail: madhavivaka360@gmail.com;

Findings showed that 30x10cm spacing ,Maximum plant height (124.48 cm), dry weight (29.85 g), number of capsules/plant (78.69), number of seeds/capsule (36.39), and seed yield (552.38 kg/ha) were obtained with 5% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*).

Keywords: Spacing; Farm Yard Manure (FYM); Recommended Dose of Nitrogen (RDN); *azotobacter*; *zaid*.

1. INTRODUCTION

Sesame (*Sesamum indicum* L.), a key oilseed crop, is produced extensively in India. As well as sesame, this seed is also known as til, simsim, benised, gingelly, and gergelim. Sesame has a key role in the agricultural and industrial industries of our country [1-4]. The only crop that produces more edible oil than sesame is groundnut. Because to the exceptional oil quality, flavour, and softness of sesame, it is referred to as "the queen of the oilseed crops" [5-7]. It typically has an oil content of 46% to 52%. By enabling optimal light contact and satisfactory absorption of nutrients and water from the soil, proper spacing in the case of sesame led to a better crop output. Sesame (*Sesamum indicum*) is a seasonal shortday plant (*Sesamum indicum* L.) [8,9]. The oldest indigenous oilseed crop in India with the longest history of cultivation is sesame (*Sesamum indicum* L.). The names til (Hindi, Punjabi, Assamese, Bengali, Marathi), tal (Gujarati), nuvvalu, manchi nuvvalu (Telugu), and ellu are all used to refer to sesame or gingelly (Tamil, Malayalam, Kannada) With 19.47 lakh acres of land and 8.66 lakh tonnes of production, tila/pitratarana (Sanskrit) and rasi (Odia) in various parts of India rank first in the globe. In comparison to other nations (535 kg/ha), India has a low average output of sesame (413 kg/ha). Sesame's low productivity is primarily due to its rainfed cultivation on marginal and submarginal fields with poor management and material shortages [10-14]. For various agro-ecological settings in the nation, better varieties and agro-production techniques are now being created. Sesame crops may produce 1200–1500 kg/ha when irrigated and 800–1000 kg/ha when rainfed with proper management. Sesame (*Sesamum indicum* L.) is one of the oldest and most significant oilseed crops, is grown around the world [15]. The plant family Pedaliaceae includes sesame. It has been cultivated in the Middle East, Asia, and Africa since 1500 BC [16]. With an average productivity of 455 kg/ha, India ranks first in both area (1.78 M ha) and production (0.81 Mt) as the world's largest producer of sesame (*Sesamum indicum* L.) It has earned the title of "the queen of

oilseeds" due to its improved nutritional, medicinal, and culinary quality [17-21]. Sesame productivity is declining as a result of its cultivation on marginal and sub-marginal and ineffective crop management techniques.

2. MATERIALS AND METHODS

The experiment was conducted in the *Zaid* of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agricultural, Technology and Sciences (SHUATS), Prayagraj (U.P.) The property is situated at 25° 57'12" N latitude and 87° 50'12" E longitude, 98 metres above mean sea level (MSL). This region is situated on the *Yamuna River's* right bank along the Prayagraj Rewa road, some 12 kilometres from the city. Ten treatments, each replicated three times and categorised as having three levels, were used in an experiment to see how spacing and nitrogen management affected the development and output of summer sesame (*Sesamum indicum* L.). The farm is situated at 25 degrees, 98 metres above mean sea level. 87° 50'12" East, 25° 57'12" North latitude; sea level (MSL).

This region is situated on the *Yamuna River's* right bank along the Prayagraj,rewa path, about 12 kilometres from the city. The experiment was set up using a randomised block design with 10 treatments that are duplicated three times in order to examine the effects of spacing and nitrogen management on summer sesame (*Sesamum indicum* L.) growth and yield.

When employed in combinations, the treatments are separated into 3 levels of plant geometry and 3 forms of nitrogen as follows: T₁: Spacing 30cm × 10cm + 50% RDN + 50% N through FYM, T₂: Spacing 30cm × 10cm + 50% RDN + Biofertilizer (*Azotobacter*) , T₃: Spacing 30cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (*Azotobacter*) , T₄: Spacing 40cm × 10cm + 50% RDN + 50% N through FYM, T₅: Spacing 40cm × 10cm + 50% RDN + Biofertilizer (*Azotobacter*), T₆: Spacing 40cm × 10cm + 50% RDN + 50% N

through FYM + Biofertiizer (*Azotobacter*), T₇: Spacing 50cm × 10cm + 50% RDN + 50% N through FYM, T₈: Spacing 50cm × 10cm + 50% RDN + Biofertiizer (*Azotobacter*) , T₉: Spacing 50cm × 10cm + 50% RDN 50% N through FYM + Biofertiizer(*Azotobacter*), T₁₀ : 10.Spacing 30cm × 10 cm + 50 : 40 : 30 NPK Kg/ha (Control) . When the oil harvest was ready, it was carefully handled. After harvesting, seeds were extracted from each net plot and dried for three days in the sun. Five randomly selected sample plants from each plot of each replication were manually measured for growth characteristics such plant height (cm), dry matter accumulation (g/plant), and more.

After cleaning and winnowing, grain yield per hectare was estimated and expressed in kilogrammes per hectare. Each net plot's leftover yield was measured and expressed,after full drying in the sun for 10 days, in tones per acre.

The statistical method developed by Gomez and Gomez was used to calculate and analyse the data (1984). The benefit:cost ratio was established after figuring out how much the seed with stover was worth, and the full cost of crop production was taken into consideration.

3. RESULTS AND DISSCUSION

3.1 Effect of Spacing and Nitrogen Management on Growth Parameters

3.1.1 Plant height (cm)

Table 1 makes it quite evident that plant height measurements increased as the crop grew.

The treatment T3 (Spacing 30cm 10cm + 50% RDN + 50% N through FYM + Biofertiizer (*Azotobacter*)) resulted in a maximum plant height of 124.48 cm, whereas the treatments T2 and T6 had statistically comparable results for spacing 30cm 10cm + 50% RDN + 50% N through FYM and Biofertiizer (*Azotobacter*).Due to nitrogen and spacing, sesame plants grew substantially higher. The spacing reveals the sesame's larger plant height, greater quantity of seeds and capsules, greater number of plants and capsules, and greatest test weight.

Solar radiation is absorbed by plants for a variety of reasons, including improved

mineral, nutrient, water, and solar radiation use, as well as plants' predisposition to extend towards light when incoming solar radiation is inadequate notably the lower canopies of plants.

The findings are consistent with those made public by Yadav et al. [22]. Because it is a crucial component of many different types of metabolically active substances, including amino acids, proteins, nucleic acids, purines and pyrimidines, nucleotides, enzymes, coenzymes, and alkaloids, nitrogen plays a significant role in plant metabolism. Increased meristematic activity of plant tissues and an increase in cell size and quantity may have contributed to higher levels of organic manure's continuous availability of nitrogen, which ultimately led to the development of taller plants. Rao et al., as well as Ghose and Sen [23], also produced results that were comparable (1990). (2000) Ghosh reported that Sesame plant height was boosted by azotobacter seed inoculation.

3.1.2 Dry matter accumulation (g/plant)

The most plant dry weight was found in treatment T3 (Spacing 30cm 10cm + 50% RDN + 50% N through FYM + Biofertiizer (*Azotobacter*)). Yet it was found that therapy T3 (spacing 30cm 10cm + 50% RDN + 50% N through FYM + biofertiizer) was statistically equal to treatment (*Azotobacter*) spacing 30 cm 10cm + 50% RDN + 50% N through FYM (*Azotobacter*). The dry weight of sesame increased significantly as a result of nitrogen and spacing. Spacing suggests that better mineral, food, water, solar radiation, etc. intake may be the cause of the increased dry weight. Nitrogen is involved in a variety of physiological processes, such as protein synthesis and enzyme activation. Reduced plant density at a 45 cm by 10 cm spacing might result in an increase in plant dryness. Plants might create weight. If they had access to enough food, water, and room. These results support Patra and Mishra's conclusions [24]. Since it improves plant photosynthesis when handled appropriately, nitrogen is an essential part of chlorophyll. Higher nitrogen levels have been found to increase the dry matter in plants because increased photosynthetic activity leads to the production of more photosynthate. Moreover, Ogundare et al. [25]. found that raising nitrogen doses to the necessary level led to improved dry matter accumulation in sesame plants.

Table 1. Effect of spacing and nitrogen management on growth attributes of Sesame

Treatment combinations	Growth attributes	
	Plant height (cm)	Dry weight (g/plant)
1. Spacing 30cm × 10cm + 50% RDN + 50% N through FYM	111.26	25.80
2. Spacing 30cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	122.89	28.72
3. Spacing 30cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)	124.48	29.85
4. Spacing 40cm × 10cm + 50% RDN + 50%N through FYM	107.65	24.38
5. Spacing 40cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	113.65	27.32
6. Spacing 40cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)	118.51	27.97
7. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM	102.56	21.36
8. Spacing 50cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	104.36	23.49
9. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)	109.86	25.49
10. Spacing 30cm × 10 cm + 50 : 40 : 30 NPK Kg/ha (Control)	98.69	20.67
F test	S	S
SEm(±)	1.55	0.45
CD (P=0.05)	4.62	1.34

Table 2. Effect of spacing and nitrogen management on yield attributes of Sesame

Treatment combinations	Yield and yield attributes					
	No.of capsules/ Plant	No. of seeds/ capsule	Test Weight (g)	Seed Yield (kg/ha)	Stover Yield (kg/ha)	Harvest Index (%)
1. Spacing 30cm × 10cm + 50% RDN + 50% N through FYM	70.89	32.72	3.34	409.65	1920.6	15.03
2. Spacing 30cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	75.61	35.28	3.38	451.64	1980	14.86
3. Spacing 30cm × 10cm + (50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>))	78.69	36.39	3.38	552.38	2052.3	17.51
4. Spacing 40cm × 10cm + 50% RDN + 50% N through FYM	68.93	29.68	3.31	401.82	2250.6	15.15
5. Spacing 40cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	72.54	34.12	3.34	438.69	2271	14.92
6. Spacing 40cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)	74.69	34.54	3.35	443.26	2501.9	14.76
7. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM	66.73	27.86	3.30	396.89	2560.8	16.70
8. Spacing 50cm × 10cm + 50% RDN + Biofertilizer (<i>Azotobacter</i>)	68.61	28.73	3.30	399.51	2315.6	16.29
9. Spacing 50cm × 10cm + 50% RDN + 50% N through FYM + Biofertilizer (<i>Azotobacter</i>)	69.76	31.33	3.32	419.36	2587.9	15.59
10. Spacing 30cm × 10 cm + 50: 40: 30 NPK Kg/ha (Control)	65.35	25.56	3.29	387.69	2602.6	16.80
F test	S	S	NS	S	S	NS
SEm(±)	0.81	0.48	0.04	8.10	2.60	0.28
CD (P=0.05)	2.40	1.43	-	24.06	7.72	-

3.2 Effect of Spacing and Nitrogen Management on Yield and Yield Attributes

3.2.1 Number of capsules/ plants

Treatment's statistical analysis of the amount of capsules produced by each plant identified a substantial impact. At a spacing of 30 cm 10 cm + (50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)), the plant produced the largest number of capsules per unit of time (78.69). Nonetheless, it was shown that the spacings of 30cm 10cm + (50% RDN + 50% N through FYM + Biofertilizer) and 40cm 10cm + (50% RDN + 50% N through FYM + Biofertilizer) were statistically equal to one another (*Azotobacter*). A sufficient quantity of sunlight absorption promotes efficient photosynthesis, which causes more photosynthates to accumulate across a larger region. With restricted spacing and a dense plant population, lower yield attribute values were attained.

3.2.2 Number of seeds/ capsules

Seed yield (kg/ha)

Notably, treatment T3 (spacing 30 cm 10 cm + 50% RDN + 50% N via FYM + biofertilizer (*Azotobacter*)) outperformed the other treatments and had the maximum number of capsules per plant (36.39). Both the T2 treatment (spacing 30 cm 10 cm + 50% RDN + biofertilizer (*Azotobacter*)) and the T8 treatment (spacing 50 cm 10 cm + 50% RDN + biofer (*Azotobacter*)). These results are in line with those of Yadav and colleagues (2007). Since it improves plant photosynthesis when handled appropriately, nitrogen is an essential part of chlorophyll. Higher nitrogen concentrations have been shown to increase dry matter because increased photosynthetic activity leads to the production of more photosynthate. In sesame, Shinde et al. [26] discovered that increasing the allowed quantity of nitrogen led to the plant accumulating dry matter. The grain yield showed that treatment T3 (spacing 30 cm 10 cm + 50% RDN + 50% N via FYM + Biofertilizer (*Azotobacter*)) outperformed the other treatments and provided the maximum quantity of seed yield (552.38 kg/ha). While it was less successful than the highest treatment T3 (Spacing 30cm 10cm + 50% RDN + 50% N via FYM + Biofertilizer), treatment T2 (Spacing 30cm 10cm + 50% RDN + Biofertilizer (*Azotobacter*)). Perhaps less

competition exists for nutrients, moisture, and light.

Adequate sunlight absorption promotes efficient photosynthesis, which causes more photosynthates to accumulate across a larger region. With restricted spacing and a dense plant population, lower yield attribute values were attained. Ogundare [25], Patra, and Mishra found findings that were comparable to these (2000).

Stover yield (kg/ha)

The yield of sesame stover was also influenced by the usage of spacing and nitrogen management. The treatment T10 (Control with specified spacing and RDN) produced the greatest recorded stover yield (2052.3 kg/ha), whereas the treatment T1 (Spacing 30 cm 10 cm + 50% RDN + 50% N via FYM) produced the lowest reported yield (1920.6 kg/ha). Perhaps less competition exists for nutrients, moisture, and light. A sufficient quantity of sunlight absorption promotes efficient photosynthesis, which causes more photosynthesis to accumulate across a larger region. With restricted spacing and a dense plant population, lower yield attribute values were attained. Potassium is involved in a variety of physiological processes, such as protein synthesis and enzyme activation. Similar findings were reported by Nayek et al. [27] and Preeti [28].

4. CONCLUSION

It can be concluded that sesame should be sown at a spacing of 30cm (row to row) × 10cm (plant to plant) and 50% Recommended dose of nitrogen through inorganic should be met out through 50% Nitrogen through FYM along with *azotobacter* seed inoculation was found more productive (552.38 kg/ha) as well as economically viable (30,221.00INR/ha).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Eshanullah, Akbar N, Iqbal A, Mohsin MR. Effect of different planting patterns on the growth and yield of two varieties of sesame. Pak J Agric Sci. 2007;44(4).
2. Ganvit JB, Sharma S, Surve VH, Ganvit VC. Effect of sowing dates and crop

- spacing on growth, yield and quality of linseed under south Gujarat condition. J Pharmacogn Phytochem. 2019;8(1):388-92.
3. Ghosh AK, Duary B, Ghosh DC. Nutrient management in summer sesame (*Sesamum indicum* L.) and its residual effect on black gram (*Vigna mungo* L.). Int J Bio-Resource Stress Manag. 2013; 4(4):541-6.
 4. Gora R, Kumar S, Singh PK, Pooniyan S, Choudhary D, PS, et al. Effect of integrated nutrient management on productivity and profitability of mustard (*Brassica juncea* L.). The Pharma. Innov J. 2022;11(2):2239-42.
 5. Alwal MA, Aktar L. Effect of row spacing on the growth and yield of peanut (*Arachis hypogaea* L.) stands. Int J Agric For Fish. 2015;3(1):7-11.
 6. Anusha O, Reddy GK, Sumathi V, Reddy PV, Reddy APK. Organic production for sustained productivity of Rabi groundnut. Andhrapradesh J Agric Sci. 2018;4(1): 56-61.
 7. Arif M, Shehzad MA, Mushtaq S. Inter and intra row spacing on growth, seed yield and oil contents of white mustard (*Sinapis alba* L.) under rainfed conditions. Pak J Agric Sci. 2012;49(1):21-5.
 8. Haritha G, Singh R, Indu T. Studies on growth and yield of sesame (*Sesamum indicum* L.) as influenced by spacing and foliar application of zinc. The Pharma. Innov J. 2022;11(3):2366-9.
 9. Helen JL, Rene G, J, Gerry G. Nature farming manual, resource efficient agricultural production-Canada. 2006;1-36.
 10. Babar S, Dongale JH. Soil fertility status as influenced by integrated nutrient management in mustard-cowpea-rice cropping sequence in lateritic soils of Konkan. Asian J Soil Sci. 2011;6(1):33-8.
 11. Beenish O, Ahmad L, Hussain A, Lal EP. Organic manure & biofertilizers: Effect on the growth and yield of Indian mustard (*Brassica juncea* L.) varieties. Curr J Appl Sci Technol. 2018;30(4):1-7. DOI: 10.9734/CJAST/2018/43594
 12. Bedigian D, Vander Maesen LJG. Slimy leaves and oily seeds; distribution and use of sesamum spp. and *Ceratotheca sesamoides* (Pedaliaceae) in Africa [international workshop]. Nairobi: Prota Foundation Wageningen. The Netherlands. 2003;271-4.
 13. Bouajila K, Sanaa M. Effects of organic amendments on soil physico- chemical and biological properties. J Mater Environ Sci. 2011;2(1):485-49.
 14. Brar ZS, Bal DS, John AS. Influences of sowing dates, nitrogen and planting geometry on the performance of Gobhi Saron (*Brassica napus* spp. olifera var. annua). Indian J Agron. 1998;43(1):133-7.
 15. Abou-Gharbia H, et al. Effects of processing on oxidative stability of sesame oil extracted from intact and dehulled seeds J. Am. Oil Chem. Soc. 1997;74: 215-221
 16. Ali GM, Yasumoto S, Seki-Katsuta M, et al. Assesment of genetic diversity in sesame (*Sesamum indicum*). Detected by amplified fragment length polymorphism markers Electronic J. Biotechnology. 2007; 10(1):0717-3458.
 17. Chandrasekaran R, Somasundaram E, Amanullah MM, Thirukumaran K, Sathyamoorthi K. Influence of varieties and plant spacing on the growth and yield of confectionery groundnut (*Arachis hypogaea* L.). Res J Agric Biol Sci. 2007;3(5):525-8.
 18. Chakraborty PC, Bairagya MD, Sarkar S, Sahoo BK, Santra GH, Nayak N et al. Effects of irrigation and nutrient management on summer sesame (*Sesamum indicum* L.). Int J Curr Microbiol Appl Sci. 2021;10(10):212-20. DOI: 10.20546/ijcmas.2021.1010.025
 19. Chetan KK, Sagar M, Tanmoy S, Monalisha P, Lalchetti S. Growth and productivity of sesame (*Sesamum indicum* L.) as influenced by spacing and nitrogen levels. Crop Res. 2022;57(3):190-4.
 20. Dalei BB, Panda S, Kheroar S, Deshmukh MR. Effect of Integrated nutrient management on growth, yield and economics of niger (*Guizotia abyssinica* L.). J Oilseeds Res. 2014;31(1):46-8.
 21. Kumar DK, Ardeshna RB, Verma BR, Patel AK. Growth and yield performance of summer Sesamum based intercropping systems. Int J Curr Microbiol Appl Sci. 2017;6(6):3341-8. DOI: 10.20546/ijcmas.2017.606.392
 22. Patil SG, Leva RL, Patil HR, Chaudhari NN. Effect of spacing and nutrient management on summer sesame (*sesamum indicum*) under south Gujarat conditions. Indian J Agricultural sciences. 2017;88(4):00-70867

23. Ghose DC, Sen JC. Analysis of yield components of sesamum as influenced by nitrogen and phosphorus fertilization. Madras Agric. J. 1980;67(10):9.
24. Patra AK, Mishra A. Effect of variety, nitrogen and spacing on yield attributes and yields of sesame (*Sesamum indicum* L.) during post-rainy season. J. Oilseeds Res. 2000;17(1):113-116.
25. Ogundare SK, Aydele FG, Oloniha JA. Effect of time of sowing and urea application rates on the growth and yield of two varieties of sesame (*Sesamum indicum* L.) in Ejiba Kogi state, Nigeria. Nigerian Journal of Agricultural, food and Environment. 2015;11(4):118-123.
26. Shinde SD, Raskar BS, Tamboili BD. Effect of spacing and sulphur levels on productivity of sesame (*Sesamum indicum* L.) under summer condition. Journal of Maharashtra Agriculture University. 2011; 36(1): 28-31
27. Nayek SS, Brahmachari K, Choudhary MR. Integrated approach in nutrient management of sesame with special reference to its yield, quality and nutrient uptake. The Bioscan. 2014;9(1): 101-105.
28. Preeti. Development of organic nutrient management for sesame *Sesamum indicum* L. M. Sc (Ag.) Thesis; 2010.

© 2023 Madhavi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/97160>