



Post-Traumatic Occlusion of the Renal Artery: A Case Report

A. Zerhouni ^{a*}, D. Farjia ^a, A. Ellouadghiri ^a, K. Khaleq ^a,
A. Bouhourri ^a and R. Harrar ^a

^a *Department of Surgical Resuscitation, Ibn Rochd University Hospital of Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98873>

Case Report

Received: 18/02/2023

Accepted: 22/04/2023

Published: 11/05/2023

ABSTRACT

The present case report highlights about Post-traumatic occlusion of the renal artery. Renal artery thrombosis is a rare complication of blunt abdominal trauma, described for the first time by Von Recklinghausen in 1861. Our case is of a 32-year-old male admitted to the emergency department to manage a polytrauma due to a car accident occurring three hours before admission. The higher mobility of the left kidney on its vascular pedicle is thought to be the cause of the increased frequency of left renal artery injuries. Because of the various therapy choices available, it is especially important to distinguish between these two conditions. The management of traumatic renal artery occlusion is controversial and opinions are divided between immediate surgical revascularization and observation.

Keywords: Post-traumatic occlusion; artery occlusion; renal artery injuries; thrombosis.

1. INTRODUCTION

Blunt abdominal trauma is routinely encountered in the emergency department. It is the main

cause of mortality and morbidity amongst the population below of the age of 35 years worldwide. Renal artery thrombosis is a rare complication of blunt abdominal trauma,

*Corresponding author: E-mail: zerhouni.amine93@yahoo.com;

described for the first time by Von Recklinghausen in 1861, however, few cases have been reported in the literature (less than 400 cases). "Pathophysiologically, this pathology can result from either intimal injury (inelastic) or compression of the renal artery against the spine by an impact on the anterior abdominal wall" [1-3]. The therapeutic management of this pathology is still a subject of controversy. In view of the particularly rare complication, the authors allowed themselves to report a clinical case of post-traumatic thrombosis of the renal artery.

2. CASE PRESENTATION

Our case is of a 32-year-old male admitted to the emergency department to manage a polytrauma due to a car accident occurring three hours before admission. The clinical examination found a patient conscious GCS 15/15 isochorous pupils, without deficit or notion of convulsions, hemodynamically and respiratory stable, he also had an epigastric and left lumbar pain. The temperature was 37.3 ° C, a blood glucose level of 1.23 G/ L. A urine test strip was made, showing traces of blood. Our initial conditioning was based on effective pain management. Laboratory investigations revealed an hemoglobin level of 9.0g/dl, WBC $13 \times 10^3 \mu\text{L}$ and platelets $88 \times 10^3 \mu\text{L}$. His blood urea nitrogen was 5.16 mmol/L, serum creatinine was 63,5 $\mu\text{mol/L}$. Lipasemia was increased to 902 IU /L. A radiological assessment was therefore launched, including normal cerebral, cervical and thoracic CT-scan, whereas abdominopelvic Contrast-enhanced CT-scan showed pancreatic contusion with hematoma and ischemic involvement of the lateral-external portion of the left kidney.

To complete the assessment a CT angiogram of the abdominal aorta was requested having shown total thrombosis of the accessory left renal artery (Fig. 1). Our management afterwards included a treatment of pancreatitis, and after multidisciplinary consultation we opted for respect of the vascular lesion.

3. DISCUSSION

A very uncommon side effect of forceful abdominal trauma is renal artery damage. However, in situations where the renal artery occlusion was thought to acceleration/ deceleration injury to the renal artery and subsequent intimal tearing, exercise and sports activities have been linked to the condition. People hurt in auto accidents usually experience

several traumas. "Polytrauma is described as two or more physical injuries to organ systems or locations, one of which may be life-threatening, leading to functional invalidity and physical, cognitive, psychological, or psychosocial impairments" [4]. Arterial spasm and arterial occlusion are the two kinds of subsequent traumatic renal vascular damage.




Fig. 1. CT angiogram of the abdominal aorta showing total thrombosis of the accessory left renal artery

The secondary effect of contusion is thought to be arterial spasm.

This syndrome shouldn't be confused with traumatic renal artery obstruction brought on by thrombosis or intimal flap formation, which might result in kidney devascularization. According to tests, the arteries' intima is the least elastic tissue and tears first when tension is applied. Due to the maximum angulation and traction on the artery at the site of fixation close to the aorta, many researchers have stated that the injury occurs inside the renal arteries 1 to 2 cm from the aorta.

The higher mobility of the left kidney on its vascular pedicle is thought to be the cause of the increased frequency of left renal artery injuries.

Because of the various therapy choices available, it is especially important to distinguish between these two conditions.

The most serious type of renal damage is devascularization of the entire kidney as a result of vascular laceration or renal artery thrombosis

(grade 5). Extensive retroperitoneal bleeding and hematuria may not be seen if the kidney is devascularized as a result of a single intimal renal artery damage that causes thrombosis.

As animal investigations have shown that kidney damage is permanent after 90 minutes of complete ischemia [5], prompt diagnosis is crucial. There aren't many particular signs or symptoms that indicate renal artery occlusion.

Reviewing 65 cases of unilateral and bilateral traumatic renal artery thrombosis from 1980 revealed that vomiting was common [6], and flank and epigastric pain were frequently observed.

All 65 patients had proteinuria upon urinalysis. Thirty percent of the patients had gross hematuria, and forty-three percent had microscopic hematuria. In 27% of cases, there was no hematuria noted.

In 1976 [7], it was shown that among 21 patients with unilateral and bilateral traumatic renal artery thrombosis, 24 percent did not have hematuria. "Radiological imaging may be helpful in detecting renal artery pathology, such as renal artery thrombosis" [8]. "The usual CT features of traumatic renal infarction include abrupt truncation/irregularity of the renal artery lumen at the point of occlusion with contrast extravasation, retrograde opacification of the renal vein from the inferior vena cava, and aberrant enhancement of the damaged kidney" [8].

Alternate linear hypodensities to delayed or reduced enhancement of the afflicted kidney are all examples of abnormal renal enhancement.

When the degree of injury grows, this "asymmetric nephrogram" becomes more noticeable, suggesting the severity of renal parenchymal ischemia [9].

The first step in treatment is to start heparin-based anticoagulation.

Enoxaparin therapy may not be necessary given the prospect of interventional radiology intervention with local thrombolysis.

This is true particularly when there is underlying renal impairment.

Anticoagulation therapy and continuing management require close collaboration between the interventionalist and the nephrologist [10]. If

there has been considerable blunt abdominal trauma and there is hemodynamic instability, surgical intervention should be pursued.

4. CONCLUSION

The treatment of traumatic renal artery occlusion is debatable, with opinions divided between rapid surgical revascularization and monitoring. This controversy has been complicated by several factors: First, the experience of each center with this type of injury is limited because of the rarity of this disease. Second, immediate revascularization has rarely resulted in a return to normal renal function, although surgeons report that the operation was technically effective at the time of the operation. Third, prolonged ischemic time due to delay in diagnosis, if beyond 12 hours, the chances of recovering renal function are gloomy.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Grablowsky OM, Weichert RF, Goff JB, Schlegel JU. Renal artery thrombosis following blunt trauma: Report of four cases. *Surgery*. 1970;67(6):895-900.
2. Laçi I, Spahiu A. Endovascular treatment of post-traumatic renal artery dissection. *Albanian Journal of Trauma and Emergency Surgery*. 2022;6(2):1086-90.
3. Khan K, Bunajem F, Alkhan F. Post traumatic arterial occlusive mesenteric ischemia: A rare case report. *Radiology Case Reports*. 2022;17(3):473-6.
4. Polytrauma/TBI system of care. US Department of veteran affairs. Available:<http://www.polytrauma.va.gov/news-and-resources/terminologyand-definitions.asp>

5. McAninch JW. Acute renal artery thrombosis following blunt trauma. *Urology*. 1975;6:74-77.
6. Barlow B, Gandhi R. Renal artery thrombosis following blunt trauma. *Journal of Trauma and Acute Care Surgery*. 1980;20(7):614-7.
7. Stables DP, Fouche RF, et al. Traumatic renal artery occlusion: 21 cases. *J Urol*. 1976;115:229-233.
8. Pyo Kang, et al. Renal infarction resulting from traumatic renal artery dissection. *The Korean Journal of Internal Medicine*. 2008;23:103-105.
9. Dobrilovic N, Bennett S, et al. Traumatic renal artery dissection identified with dynamic helical computed tomography. *J Vasc Surg*. 2001;34:562-64.
10. Sauerberg N, Khan YS. Renal artery thrombosis. In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2023. Available: <https://www.ncbi.nlm.nih.gov/books/NBK564380/>

© 2023 Zerhouni et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98873>