



A Case Study on Seasonal Behaviour of *Aphodius elegans* Allibert an Endocoprid Dung Beetle

Nythri Tshering ^{a*}

^a Department of Environment & Life Sciences, Sherubtse College, Royal University of Bhutan, Bhutan.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajriz/2024/v7i3156>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:
<https://www.sdiarticle5.com/review-history/117918>

Received: 12/04/2024

Accepted: 15/06/2024

Published: 19/06/2024

Original Research Article

ABSTRACT

Dung beetles (Scarabaeidae) classified into endocoprid, paracoprid, and telecoprid beetles, are fascinating insects with various behaviors. The study focuses on behaviour of endocoprid beetles *Aphodius elegans* Allibert, found in Bumdeling, Trashi Yangts Bhutan. The observational and scan sampling methods were employed during the study. These beetles locate dung using antennae and dig inside it, resembling a boat propelled by oars. They are active during warm weather and bury themselves under dung during cold. Beetles defecate semi solid feces after dung foraging. They mimic death as defensive and antipredator behaviour. They prefer moist dung pats for burial and use repeated attempts and strongholds of hindwings to initiate flight. This study substantiates the diverse fascinating behaviors of a less-focused group of insects to gain a deeper understanding of their interactions with ecology.

Keywords: *Aphodius elegans* Allibert; behavior, dung beetle, endocoprid.

*Corresponding author: Email: nythripenjor@gmail.com;

Cite as: Tshering, Nythri. 2024. "A Case Study on Seasonal Behaviour of *Aphodius Elegans Allibert* an Endocoprid Dung Beetle". *Asian Journal of Research in Zoology* 7 (3):53-59. <https://doi.org/10.9734/ajriz/2024/v7i3156>.

1. INTRODUCTION

Dung beetles are insects of order Coleoptera, consisting the species of the subfamilies; Scarabaeinae, Geotrupinae, and Aphodiinae [1] that partly or exclusively feeds on feces throughout their lifecycle [2,1]. They are found in every continent except Antarctica. Africa is considered to have the most diverse population of dung beetles [3]. Historically, since ancient Egyptian, dung beetles were believed as the representation of rebirth and rejuvenation [4]. They were revered as the symbol of the god of the sun, who rolled the sun across the sky each day and buried it in the evening, only to emerge again the next morning [3].

Ecologically, dung beetles are a major component of biological dung removal [5] and controllers of pests and parasites that breed in dung [6,5]. Dung beetles help farmers in agriculture production, by reducing the yield loss for pests, cost of pesticide applications, and fertilizer use. They are considered bioengineers of the environment and health giver for the population [3]. Besides their role as decomposers, dung beetles maintain biodiversity through its involvement in the Pollination and seed dispersal of some plants [7,8].

Dung beetles are adapted to a wide variety of ecosystems and many are highly specialized [4]. Many specialist dung beetle species can feed on resources such as leaves, fungus, rotten fruits and small insects other than dung [2]. Depending upon the species, dung beetles have a specific preference towards dung, dung condition, and the dung odor [9]. Dung beetles can be segregated based on habitat and soil type. The activity and the behavior exhibition of dung beetles varies with species, the season and the seasonal activity, and the light and light intensity [4]. Additionally, beetles respond differently to elevation and temperatures [3].

Most dung beetles are generalist (coprophagous) feeders, however reduced dung availability sometime leads to specialization in dung beetles [4]. Along with abiotic elements as weather and environment, feeding habits are also impacted by predatory and competitive variables [3]. Dung beetles have detritus-feeding habits and a dung resource contains all the nutrition that the beetles require. Adults have filtering mouth parts that feed on a “dung Slurpie”, the primary moisture within the dung [4]. While larvae have biting mouthparts, that makes it possible to feed on

both moisture and the fiber of the dung within the brood balls [3].

Dung beetles are divided into three behavioral groups; rollers, tunnelers, and dwellers. The beetles that roll dung into round balls, used for food sources or brooding chambers are rollers. The tunnelers make a tunnel and bury the dung wherever they find it. The dwellers neither roll nor burrow the dung, but they simply live-in dung [4]. The dung beetles are also known as Endocoprid (dwellers), Paracoprid (tunnelers) or Telecoprid (rollers) insects [10,4,5].

The fascinating behavior of the two groups of the dung beetle, the roller and tunneler were widely studied. Rollers form a bit of dung ball, roll it away and bury it. The dung balls are used either by females for laying the egg as a brood ball or as a food source for adults [4,2]. The male and female rollers establish a pair bond after encountering the dung pat. The male prepares the giant dung ball and offers it to the female as the brood ball. If she accepts, then they roll it away together or the female beetle rides on the top of the ball [4]. They cooperatively defend their ball from stealing by other beetles [11]. The warfare and the battle between the males are fought to show strength and bravery [10]. The new pair bond finds a suitable and safe place to bury the ball before mating. The male leaves the scene, while the female lays a single egg and prepares another brood ball. The brood ball is then coated and sealed with a mixture of dung and saliva. Some mothers stay with the ball till the larvae hatch [4,11].

The tunnelers dive into the dung and tunnel directly into the soil to start the family [12,4]. Storing the dung underground helps to protect the developing larvae from predators and also to keep the nutrients, in the dung pile fresh and digestible [2]. The female arranges and sorts the dung in the tunnel while the male carries the fresh dung to the home. The parents stay with the larvae until they are matured [2,13]. The dweller neither rolls nor makes tunnels. The female lays eggs on the top of the dung and the larval development takes place inside the dung pat [3].

The behavior of the dweller beetles was not exclusively studied as it has no fascinating behavior like that of roller and tunnelers. Therefore, the study focuses on the different behaviors shown by the dweller dung beetle *Aphodius elegans* Allibert. The different

behaviors performed was observed and explained accordingly. The behaviors of hunting the dung pat, shoveling the dung pat and their burial, their feeding habits, the defecation of feces, and also their defensive behaviors were explained in the study.

2. METHODOLOGY

2.1 Study Site

The dung beetles were collected from the small cow herds of the Betsamang village, Bumdeling gewog (village block) under Trashigang Dzongkhag (district). The small study site was created by a collection of fresh dung pats of the cow. The dung inhabited with the *Aphodius elegans* Allibert was also collected and kept in the study site. The study was carried out during the spring season (March and April) of 2020. The region falls under Bumdeling Wildlife Sanctuary at elevations between 1500 meters (4,900 ft) and 6000 meters (20,000 ft). The mean annual summer temperature is 22.6 °C and the mean annual winter temperature is 15 °C.

2.2 Materials

Fresh dung pat, a spade, Gloves, Camera (Mobile camera).

2.3 Sample Collection

The fresh cow dung and *Aphodius elegans* beetles were collected at a specific site for observation. Field observational [14], and scan sampling methods [15] were used, and videos of their behaviors were recorded. Observed behaviours were described and verified with literatures available. Dung beetles were counted and identified, and then released in an open area.

2.4 Procedure to Hunt Dung Beetles

Examine cow herds and pastures for dung pats, looking for holes or shredded surfaces. Look for adult beetles and larvae. Observe insect activity immediately after dung deposition. Dung beetles arrive within minutes at temperatures above 70°F [12]. *Aphodius elegans* beetles appears twice a year in spring and summer [16].




Fig. 1. Study site; Bumdeling, Trashiyangtse, Bhutan

3. RESULTS AND DISCUSSION

3.1 General Body Anatomy of Dung Beetle

Body composed of 3 main segments; head, thorax and abdomen. A typical characteristic of beetles is the hard elytra that varies in color from black, brown, yellow or purple, depending on the species of dung beetle. A pair of wings folded under the elytra and hardened forewings covers the body. It possesses 3 pairs of legs terminated with appendages specialized for shoveling the dung and earth. Male of some species have a pair of horns on their head, used for defense. Antennae are typically characterized by elongated and segmented with a discernible curvature. The size of dung beetle varies from the smallest species 2mm long to the larger species up to 30mm long [17].

3.2 Taxonomy of *Aphodius elegans* Allibert

Class: Insecta

Order: Coleoptera

Superfamily: Scarabaeoidea

Family: Scarabaeidae

Subfamily: Aphodiidae

Genus: *Aphodius*

Species: *Aphodius elegans* Allibert

3.3 Identification Features of *Aphodius elegans* Allibert

The identification features of the studied dung beetle are in accordance to Italian Entomologist Forum, [18]. *Aphodius elegans* is identified by its auriculate genae with flat scutellum. Elytra interstices are nearly flat, shiny, sparsely or finely with distinct punctured. Blackish, elytra pale yellowish with bands usually widened till suture. The first segment of the hind tarsi conical, is slightly longer than the following three combined. Fore tibia, apical spur, in males outwardly hooked.

3.4 Behaviors of *Aphodius elegans* Allibert

Dung pat hunting behavior: Adult beetles locate food sources by smell and detect it during flight using antennae. They eat in situ and may be transported for further consumption [10]. Adult

Aphodius elegans use antennae to locate dung pats and feed on liquid content [19]. Beetles actively move towards freshly scattered cow dung, occupying it on the second day of observation.

Aphodius elegans migrate to new dung pats in summer, depending on weather and dung condition. Dormont et al., [9], mentioned that different species have specific preferences of dung with dung condition and dung odor. *Aphodius elegans* leave dung pats when it becomes hard and dry, leaving earlier inhabited ones 9 days after first observation. They rely on scattered dung for food.

Dung excavation and shoveling behavior: Adult dung beetles orient themselves towards the dung surface, quickly burying themselves under it. They prefer the fresh or soft part of the dung for their burial. They use antennae to detect food sources and examine the dung pat for entrance [19]. Beetles set their antennae in motion and explore dung pats, examining fresh and semisolid parts for entrance. Their morphological adaptive features aid in their successful habitation.

The *Aphodius elegans* dung beetle buries its clypeus and head under the dung pat using protibial and mesotibial muscles. Its protibia, mesotibia, and metatibia help it enter the dung pat, clearing the path and propelling the body forward. The beetle's excavation and movement resemble boats propelled by oars, with each leg having an appendage for shoveling dung and earth. The dung beetle's legs are adapted for digging [20].

Hibernating behavior: According to Yoshida & Katakura,[21] and Finn & Gittings, [19], Adult *Aphodius elegans* are observed twice a year, in spring and autumn. The first flight occurs in July, with abundant synchrony, and the second flight occurs in September to October, containing mature eggs [21].

Aphodius elegans excavate chambers under dung pats to feed on liquid content. They retract their appendages and live motionlessly, storing energy and preventing cold. Adult dwellers are found in moist droppings and drying dung larvae. Beetles hibernate as mature larvae concentrating in early mornings and cold weather [21].




Fig. 2. Morphological feature of *Aphodius elegans* Allibert

Feeding behavior: Beetles scramble and consume dung, primarily derived from microorganisms. Their mouth part is well adapted for handling soft, pasty food dung [22,23]. According to Hata & Edmond, [23], mandibles in *Aphodius elegans* grind particulates, while molae functions as a pestle mortar system. These actions are observed in adult *Aphodius elegans* through the grinding and squeezing action of the dung beetle.

Aphodius elegans beetles have a distinct maxillary palp for food scrapping, but prefer moist, fresh dung pats [19], resulting in a "dung slurpie" feeding habits [4] making them scrapper and filter feeders.

Defecating Behavior: The *Aphodius elegans* dung beetle excretes waste and feces in its habitat, but their actions are unclear. They defecate while feeding on the dung pat, leaving dung traces of their movements on white paper. The dung beetle's path is marked with ink patterns of semi solid feces, turning the dung into fine soil.

Defensive behavior; the death mimic: *Aphodius elegans* beetles in open spaces orient themselves towards dung and find hiding places to avoid danger. Touching their bodies retards appendages and mimics death. Quick burial under dung prevents predation and environmental effects. Leg spurs and strong appendages aid fighting and digging. They prefer

dark, shaded areas for quick burial under dung pats. They quickly reposition their bodies during sunny days to avoid danger and find fresh dung pats for burial.

Body repositioning and flight start behavior: Adult *Aphodius elegans* use their tibia and appendages to regain their normal walking position when upside down or accidentally fallen from a dung pat. Their strong specialized tibia helps them attach to the ground surface.

Beetles attempt to reposition their body after initial failures using elytra and hindwings. The elytra, a hardened structure, uplifts the thorax and abdomen while pushing the head towards the ground [24]. The fore tibia is stretched forward to hold onto supports, and the behavior is repeated at regular intervals until they success.

Flight behavior: *Aphodius elegans* beetles primarily observed during bright sunlight or warm weather, exhibit active flight behavior. Beetles hold out the forewing elytra and hindwing, generating lift [24] and used the hindwing for rigorous flapping [25].

4. CONCLUSION

Dung beetles, coprophagous insects, which exhibit fascinating behaviors. However, early studies had focused mostly on dung roller and tunneler beetles, and endocorpid dweller dung

beetles require more focus. Thus, *Aphodius elegans*, endocorpid dung beetles, are used for behavioral studies in the Bumdeling region.

The adult beetle prefers moist, fresh dung pats for habitat. When near these pats, it sets its antennae to search for food. The adult *Aphodius elegans* directs its body towards the dung, shovel and buries under it. The protibia assists in quick burial, while the mesotibia and metatibia push the body forward, similar to a boat propelled by oars.

They use their specialized mouth to scrape and filter dung, using excretion to convert dung into fertile soil. They use defensive mechanisms like death mimics and retracting appendages under the hard elytra. Their elytra is used for repositioning and launching flight, with the tibia retracting to avoid sunlight and their strong outlet of wings generate the body lift.

Aphodius elegans are active during warm weather, with their diversity varying with the season. However, extensive study on mating, competition, and lifecycle of these beetles is limited due to lack of tools for detailed anatomical studies and behavior analysis. The portrayal of other fascinating behaviors may be hindered by disturbances in their activity and habitat.

Further research is needed to understand mating, competition, and warfare battles among beetles. Studying feeding and flying behaviors of *Aphodius elegans* is essential. Observing the dung beetle's behaviors twice a year, in spring and autumn, can provide insights into their hunting, excavation, shoveling, feeding habits, and flight techniques. The bright sunny days are ideal for studying these fascinating behaviors.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

I hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Huerta C, Oca EM De, Rosales MC, Favila ME. CHAPTER 24 the role of dung beetles in the sustainability of pasture and grasslands; 2013.
Available:<https://doi.org/10.2495/978-1-84564-75>
2. Holter P, Scholtz CH. What do dung beetles eat? Ecological Entomology. 2007;32:690- 697.
3. Whipple SD. Dung Beetle Ecology: Habitat and Food Preference, Hypoxia Tolerance, and Genetic Variation, Dissertations and Student Research in Entomology. 2011;12.
4. Hanski I, Cambefort Y. Dung beetle ecology. Princeton University Press. Princeton, New Jersey. 1991;481.
5. Byk A, Pietka J. Dung beetles and their role in nature; 2018.
Available:<https://doi.org/10.24131/3247.180103>.
6. Fincher GT. Effects of dung beetle activity on the number of nematode parasites acquired by grazing cattle. The Journal of Parasitology. 1975;61(4):759-762.
7. Shepherd VE, Chapman CA. Dung beetles as secondary seed dispersers: Impact on seed predation and germination. 1998; 199–215.
8. Griffiths HM, Bardgett RD, Louzada J, Barlow J, Griffiths HM. The value of trophic interactions for ecosystem function: Dung beetle communities influence seed burial and seedling recruitment in tropical forests; 2016.
9. Dormont L, Epinat G, Lumaret J. Trophic preferences mediated by olfactory cues in dung beetles colonizing cattle and horse dung. Environmental Entomology. 2004; 33(2):370- 377.
10. Halffter G, Edmonds WD. The nesting behavior of dung beetles (Scarabaeinae). An ecological and evolutive approach. Publ. 10, Instituto de Ecología AC.: México. 1982;176.
11. Baird E, Byrne MJ, Smolka J, Warrant EJ, Dacke M. The Dung Beetle Dance: An Orientation Behaviour?. 2012;7(1):16.
Available:<https://doi.org/10.1371/journal.pone.0030211>.
12. Bertone M, Watson W, Stringham M, Green J, Washburn S, Poor M, Hucks M. (n.d.). Dung Beetles of Central and Eastern North Carolina Cattle Pastures. 2004;1–7.
13. Dabrowski J, Venter G, Truter WF, Scholtz CH. Dung beetles can tunnel into highly compacted soils from reclaimed mined sites in e Malahleni, South Africa Dung beetles can tunnel into highly compacted

- soils from reclaimed mined sites in e Malahleni, South Africa. *Applied Soil Ecology*. 2018;134(November):116–119. Available:<https://doi.org/10.1016/j.apsoil.2018.10.015>.
14. Altmann J. Observational sampling methods for insect behavioural Ecolgy. *Florida Entomologist*. 1984;67(1):50–56. Available:<https://www.jstor.org/stable/3494104?seq=1>
15. Koster J, McElreath R. Multinomial analysis of behavior: Statistical methods. *Behav Ecol Sociobiol*. 2017;71(131).
16. Bang H, Han M, Kang K, Lee D. Diversity of coprophagous invertebrates in Korea and their role in soil ecosystem-mainly dung beetles. 2017;1–12.
17. Thomas JC, Darren JM. Identification guide to the Scarabaeinae dung beetles of Cusuco National Park, Honduras; 2012.
18. Italian Entomologist Forum. *Aphodius elegans Allibert, 1847-Aphodiidae-e specie simili, Review of the genus Aphodius (Coleoptera)*. Italy; 2017.
19. Finn JA, Gittings T. Review A review of competition in north temperate dung beetle communities; 2003.
20. Kerbtier BD. Beetle morphology. Christoph Benisch, Baden-Wurttemberg, Germany; 2007.
21. Yoshida, N & Katakura, H. Life Cycles of Aphodius Dung Beetles (Scarabaeidae, Coleoptera) in Sapporo, Northern Japan. *Environmental science, Hokkaido: journal of the Graduate School of Environmental Science, Hokkaido University, Sapporo*, 1986;8 (2):209-229.
22. Miller A. The mouth parts and digestive tract of adult dung beetles (Coleoptera: Scarabaeidae), with reference to the ingestion of helminth eggs. *J. Parasitol*. 1961;47:735-744.
23. Hata K, Edmond WD. Structure and Function of the mandibles of the adult dung beetles (Coleoptera: Scarabaeidae), *Internal Journal of Insect Morphology and Embryology*. 1983;12(1):1-12.
24. Westheim CJ, Engel S, Baird E, Dacke M, Muijres FT, Hedenstrom A. Beetle flight: Flapping protective wings increase the lift. *Science Daily*; 2012.
25. Fincher GT. The potential value of dung beetles in pasture ecosystems. *Journal of the Georgia Entomological Society*. 1981; 16(1):316-333.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117918>