



Distribution and Antibiotic Sensitivity Patterns of Faecal Bacteria from Underground Faecal Storage Cisterns in Ondo State, Nigeria

Anthony Kayode Onifade¹ and Sunday Stephen Abi^{1*}

¹Department of Microbiology, Federal University of Technology, Akure (FUTA), Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author AKO designed the study and supervised the work. Author SSA performed the statistical analysis, wrote the protocol and the first draft of the manuscript. Author SSA carried out the microbiological analysis of the study, with thorough supervision by author AKO. Author SSA also carried out the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMB/2020/v20i130206

Editor(s):

(1) Dr. Simone Aquino, Instituto de Pesquisas Energéticas e Nucleares (IPEN), Brazil.

Reviewers:

- (1) Bagiu Radu-Vasile, University of Medicine and Pharmacy Timisoara, Romania.
(2) Paola Taydé Vázquez Villegas, Instituto Tecnológico Superior de Cintalapa, México.
(3) Claudious Gufe, Central Veterinary Laboratory, Zimbabwe.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/54412>

Original Research Article

Received 28 November 2019

Accepted 01 February 2020

Published 08 February 2020

ABSTRACT

Two hundred and fifty-six (256) human faecal samples were collected from thirty-three (33) underground faecal storage cisterns across 5 major towns (Akure, Akungba, Ondo, Ore and Owo) in Ondo State, Nigeria. Salmonella-Shigella agar, Eosine Methylene Blue agar and MacConkey agar, which are selective for faecal bacteria, were used to isolate a total of 103 strains of Gram-negative bacteria. Biochemical characterization of the isolates revealed *Escherichia coli* (32.04%), *Enterobacter aerogenes* (14.56%), *Klebsiella pneumoniae* (12.62%), *Salmonella typhi* (11.65%), *Pseudomonas aeruginosa* (7.77%), *Shigella dysenteriae* (6.80%), *Proteus mirabilis* (5.83%), *Citrobacter koseri* (3.88%), *Providentia alcalifaciens* (2.91%) and *Klebsiella oxytoca* (1.94%). All the bacterial isolates were then subjected to antibiotic sensitivity test using antibiotic discs impregnated with Augmentin (30 µg), Gentamycin (10 µg), Pefloxacin (10 µg), Ofloxacin (5 µg), Streptomycin (10 µg), Sulfomethoxazole-trimethoprim (30 µg), Chloramphenicol (30 µg); Sparfloxacin (10 µg); Ciprofloxacin (5 µg) and Amoxicillin (20 µg). The resistance of the isolates to

*Corresponding author: E-mail: abisundaystephen@gmail.com, abisundaystephen@yahoo.com;

the antibiotics used varied from 27.18% to Ciprofloxacin, 33.98% to Ofloxacin, 47.57% to Sulphomethazole-Trimethoprim, 56.31% to Streptomycin, 66.99% to Chloramphenicol, 74.76% to Pefloxacin, 75.73% to Sparfloxacin, 75.73% to Gentamycin, 85.44% to Ampicillin and 95.15 to Augmentin. These results showed that underground human faecal storage cisterns contain potential multiple antibiotic resistant bacteria that may be transmitted to human via fomites, animal vectors and water.

Keywords: Cisterns; faecal; antibiotics; underground.

1. INTRODUCTION

Antibiotic resistance has been defined as the ability of bacteria to change in ways that resist the effects of drugs – “that is, the germs are not killed and their growth is not stopped”. In other words, antibiotic resistant bacteria are bacteria that are no longer susceptible to the antibiotics to which they were earlier susceptible. Antibiotic resistance in bacteria remains a major problem and the environments that help to maintain such resistance represent significant problem in the community [1]. During the past sixty years, antibiotics were extensively used in humans and in veterinary medicine, as well as in breeding practices. Continued high rates of antibiotic use in hospitals, the community and agriculture have contributed to selection pressure that has sustained resistant strains [2], forcing a shift to more expensive and more broad-spectrum antibiotics. Despite concerted efforts being made to control antibiotic resistance, the spread of antibiotic resistant bacteria continue to be an important global problem, particularly in developing and low income countries, where they rapidly spread, causing morbidity and mortality with an attendant increase in the cost of treating infectious diseases due to treatment failures. In fact, the recent and so far the first, global report on antimicrobial resistance by WHO indicates alarmingly high rates of resistance in the bacteria which cause common infections in healthcare facilities and in the community. Meanwhile more than 200,000 newborns are estimated to die each year due to infections for which effective antibiotics are unavailable [3]. This is becoming a global threat [4]. In fact, it is estimated that the global annual mortality of 700, 000 death associated with antibiotic resistance may by 2050 increase to 10 million, at a cost of 100 trillion USD to the world economy, if nothing is done [4]. There is a large group of Gram negative bacteria called Enterobacteriaceae. They are enteric bacteria and are normal intestinal flora of human. Although they are not pathogenic under normal conditions, they are capable of causing enteric

infections mostly in immune compromised hosts. Enteric infections are the fifth leading cause of death worldwide. Nearly 70% of such infections are food-borne. About 1.5 billion cases of diarrheal disease occur annually, killing 2.2 million people, mainly children [5]. Nigeria is also confronted with the burdens of antimicrobial resistance. The Nigerian Centre for Disease Control (NCDC), in collaboration with other institutions, has made efforts to develop an approach to combat antibiotic resistance using an evidence-based method. Meanwhile, [5] reported that Nigeria has experienced huge resistance to antimicrobials in humans, especially in sepsis, respiratory and diarrheal infections. These include childhood-related life-threatening diseases and are supported by empirical evidence, which are replete and scattered in peer-reviewed and grey literature, as well as commissioned reports. In addition, the situation analysis and recommendations on antimicrobial resistance and drug use in Nigeria has recently been documented [5]. For example, Nigeria’s diarrhoea prevalence rate is one of the worst in sub-Saharan Africa. It is 18.8% and notably higher than the continental average of 16% [6]. Failure to control the spread of diarrhoeal pathogens both resistant and non-resistant ones have greatly worsened the burden of diarrhoeal disease in Sub-Saharan Africa.

Bacteria efficiently exchange genetic materials, particularly among related species and/or groups including the pathogenic ones. And the intestine is a suitable environment for these activities and therefore intestinal flora is a reservoir for resistance bacteria genes and they are passed out with human faeces. About 70% of the antibiotics produce globally are used in agriculture, the remaining 30% are consumed by man. Significant proportions of antibiotics consumed by man are excreted and passed into the environment [4]. In communities with less developed sanitation in fracture, there is a higher risk that waste will not be treated and sometimes be closer to communities, thus increasing the risk of exposure to the carriage of resistant bacteria

by otherwise healthy people and the rate of drug-resistant community-acquired infections. It is in these settings that there is an additional concern about antibiotics and resistant bacteria passing into the environment as sewage system are not often functional. Inappropriate human disposal of antibiotics, for instance by flushing them into the toilet, plays role in this [4]. Toilet users who go into the toilet with mobile phone can leave the toilet with myriads of pathogenic bacteria. Also, by vertical and horizontal flow of faecal sludge especially during raining season, the bacteria in human underground faecal storage tanks can leak into water aquifers and wells that are not in safe distance to the faecal storage tanks. Toilets act as a vehicle for the transmission of pathogens from gut, respiratory tract and skin via hands and surfaces from one person to another [7]. Due to unhygienic use of toilet facilities, faecal matter remains a major reservoir source of human pathogens. When hands containing faecal remnant uses a door knob, the bacteria pass on to it.

Although there are numerous documented reports on antibiotic sensitivity patterns of faecal bacteria isolated from various sources, there are scanty reports on antibiotic sensitivity patterns of bacteria isolated from human excreta (mixture of faeces and urine) collected from underground faecal storage cisterns. This report was therefore designed to determine the distribution and antibiotic sensitivity patterns of faecal bacteria isolated from human excreta collected from underground faecal storage cisterns in Ondo State, Nigeria.

2. MATERIALS AND METHODS

2.1 Description of Study Location

This research was carried out in Ondo State, Nigeria from January to November, 2019. Ondo was created on 3rd February, 1976, from the former Western State. It originally included the present Ekiti State, which was split off in 1996. Akure is the administrative capital of Ondo State. The State, with coordinates 7°10'N 5°05'E, has a land area of 14, 606 km² and human population of 3,460,877; the statistic of males stood at 1,745,057 while females was 1,715,820 (2006 census). However, the projected population of Ondo State as at year 2016 was 4,671,700. The State borders: Ekiti State to the north, Kogi State to the northeast, Edo State to the east, Delta State to the southeast, Ogun State to the southwest and Osun State to the northwest.

2.2 Collection of Samples

Two hundred and fifty-six human faecal samples were aseptically collected in different screw-cap sterile bottles from thirty-three faecal storage tanks across five towns (Akungba, Owo, Akure, Ondo and Ore) in Ondo State, Nigeria, from the month of January to May 2019. All samples were transported in an icepark box within 3 hours of collection to the Microbiology Laboratory of the Federal University of Technology Akure, where bacteriological analysis was carried out on them.

2.3 Preparation and Sterilization of Agar

Nutrient agar (BioLab, USA), MacConkey agar (HiMedia, India), Eosin Methylene Blue (EMB) agar (BioLab, USA) and Salmonella-Shigella (S-S) agar (BioLab, USA) were prepared following manufacturers recommendations. All the prepared media, except Salmonella-Shigella agar, was then autoclaved at 121°C for 15minutes. Salmonella-Shigella was dissolved by heating on hot plate [8].

2.4 Enumeration and Agar-Dependent Isolation

Serial dilutions of the faecal samples were carried out aseptically up to 10⁻⁶ dilution in order to obtain countable bacteria colonies on the agar plates. Using pour plate method, 1 ml aliquot from the dilution 10⁻⁵ and 10⁻⁶ was poured on separate duplicated sterile Petri dishes containing sterile nutrient and MaConckey agar. The agar was allowed to congeal on the poured plates and was then incubated invertedly for 18-24 hours at 37°C in a bacteriological incubator. Nutrient agar was used for total bacterial count while MaConckey agar was used as selective agar for faecal bacteria. Colonies on the agal plates were counted and recorded. Dinstict colonies from the MaConckey agal plates were picked with the aid of sterilized wire loop and streaked on Salmonella-Shigella agar and Eosin Methylene Blue agar, for purification. Pure colonies were later stored at 4°C on Nutrient Agar (NA) slant in a refrigerator [9].

2.5 Biochemical Characterization

To further identify the isolates, various biochemical and sugar fermentation tests were carried out on each isolates. Gram staining, Oxidase, Urease, Catalase, Coagulase, Citrate, SIM (Sulphur, Indole and Motility), Ornithine, Lysine, Methyl Red, Voges Prauskeur, TSI

(Triple Sugar Iron) tests were carried out. Sugar fermentation (glucose, sucrose, mannitol and Xylose) was also carried out [8]

2.5.1 Gram staining

A loopful inoculum of 24-hour pure bacterial cultures was speared on grease-free glass slides. The smears were air-dried and then heat-fixed by rapidly passing it through flame emanating from burnsen burner. This is to maintain the cellular integrity of the bacterial cells and prevent it from washing away during flooding. The smears on the slides were then flooded with crystal violent dye (primary stain) for 60 seconds and then quickly washed off under a slow running clean water. Subsequently, the smears were covered with lugol's iodine (mordant) for 60 seconds and washed off with clean running water. 75% ethanol (decourlorizer) was poured on the smears and washed of after about 30 seconds and was again washed off under slow flowing tap water. Finally, Sefranin (secondary stain) was added for 60 seconds and was washed off under slow running tap water. The slides were then left to air dry, after which they were viewed under 40x objective lens of a light microscope. A drop of oil immersion was placed on the stained smear and was again examined using using oil immersion objective (x100) lens of the same microscope [10,8].

2.5.2 Oxidase test

A piece of filter paper was soaked with 2-3 drops of oxidase reagent. Using a sterile piece of stick, a colony from 24 hours culture of the test organism was picked and smeared on the filter paper; reactions was observed for 10 seconds. Oxidase positive isolates changed to blue or purple colour, while oxidase negative isolates has no colour change within 10 seconds [8].

2.5.3 Urease test

This was done by streaking overnight culture of the test organisms over the slant surface of prepared slanted urease agar in a test tube. A tube without any test organisms was used as control. Release of ammonia brings about colour change from yellow to pink or red which means a positive result; tubes with no colour change werw regarded as containing organism tested negative to urease production [10].

2.5.4 Catalase test

The test distinguished catalase producing bacteria from non-catalase producer. Catalase

positive isolates breakdown hydrogen peroxide to oxygen and water. 3% hydrogen peroxide was prepared and a drop was placed on a glass slide. Using a sterile wooden stick, a colony of the test organism from 24 hours old culture was placed in the hydrogen peroxide and mixed together gently. Copious bubbles caused by the organism by the liberation of oxygen indicated positive result [8].

2.5.5 Coagulase test

A 24 hrs old culture was emulsified in normal saline on clear grease free slide containing a drop of distilled water on both end of the slide to make two thick suspensions. A loopful of plasma was added to one of the suspensions and gently mixed. The second suspensions contain no plasma and was used as control to differentiate any granular appearing as coagulase of the organism from clumping caused by the organism's Coagulase enzyme. Clumping within 10 seconds indicated a positive result while, no clumping indicated a negative result [8].

2.5.6 Citrate test

Simon citrate agar was prepared in a beaker following manufacturer's instruction. The beaker containing the prepared agar was homongenized on electric hot plate. Using sterile syringe, about 10 ml each was dispensed into different test tubes. The component was then autoclaved at 121°C for 15 minutes. The test tubes were slanted and allowed to congeal. Overnight culture of the test organism was inoculated on the slanted portion of the medium and incubated at 37°C for 2-5 days. Colour change from green to bright blue colour indicated a positive citrate test while, no colour change indicated a negative citrate test of medium [11].

2.5.7 SIM (Sulphur, Indole, and Motility) test

SIM agar was prepared according to the manufacturer's specification in a conical flask. It was homogenized on electric hot plate before dispensing about 10 ml each into various test tubes. The components was then sterilized in the autoclave for 15 minutes at 121°C. After the agar cooled and turned semi-solid, 24 hrs cultures of the test organisms were inoculated into the tubes by making a stab on the center of the medium to a depth of about one (1) inch and then incubated aerobically at 37°C for 24 hours in a bacteriological incubator. H₂S positive test indicated by blackening of the medium along the inoculation line. H₂S negative result means there

is no blackening. Positive motility test indicate by a diffused zone of growth from inoculation line whereas a negative result was infected when there is no diffuse growth. In order to examine for indole production, 3-6 drops of Kovac reagent was added into the tubes. A reddish ring means the test organism is positive to indole production [10].

2.5.8 Triple Sugar Iron (TSI) test

TSI agar was prepared in a conical flask according to the manufacturer's specification. It was homogenized on electric hot plate before dispensing about 10 ml each into test tubes. The tubes were then sterilized in the autoclave for 15 minutes at 121°C. After autoclaving, the test tubes containing the TSI was slanted and left to congeal. Using aseptic technique, colony from overnight pure bacterial culture was picked with a sterile straight needle and stabbed into the medium and then streaked the needle back and forth along the surface of the slant. The neck of the TSI tube was cocked and flamed before incubating at 37°C for 18 to 24 hours. Alkaline slant (red) and acid butt (yellow) with or without gas production (breaks in the agar butt) indicated that glucose fermentation has occurred. The organisms preferentially degrade glucose first. Since this substrate was present in minimal concentration, the small amount of the acid produced on the slant surface was oxidized rapidly. The peptones in the medium was also used in the production of alkali. At the butt, the acid reaction is maintained because of the reduced oxygen tension and slower growth of the organisms [8].

Acid slant (yellow) and acid butt (yellow) with or without gas production indicated that lactose or sucrose fermentation has occurred. Since these substances are present in higher concentrations, they serve as substrates for continued fermentative activities with maintenance of an acid reaction in both the slant and the butt. Since these substances are present in higher concentrations, they serve as substrates for continued fermentative activities with maintenance of an acid reaction in both the slant and the butt.

Alkaline slant (red) and alkaline butt (red) or no change (orange-red) butt indicated that no carbohydrate fermentation has occurred. Instead; peptones was catabolized under anaerobic and /or aerobic conditions resulting in alkaline pH due to production of ammonia [8]. The presence of

hydrogen sulphide was indicated by blackening of the TSI agar. Some bacteria utilize thio sulfate anion as a terminal electron acceptor, reducing it to sulfide. If this occurs, the newly-formed hydrogen sulfide (H_2S) reacts with ferrous sulfate in the medium to form ferrous sulfide, which is visible as a black precipitate [8].

2.5.9 Ornithine decarboxylation test

Decarboxylation is the process by which bacteria that possess specific decarboxylase enzyme attack amino acids at their carboxyl end (-COOH) to yield an amine or a diamine and carbon dioxide. The amino acid L-ornithine is decarboxylated by the enzyme ornithine decarboxylase to yield the diamine putrescine and carbon dioxide [12]. Ornithine decarboxylase broth was used for this test. The broth contains L-Ornithine monohydrochloride, Yeast extract, Glucose and Bromo cresol purple. Yeast extract in the medium provides nitrogen and other nutrients necessary to support bacterial growth. The amino acid ornithine is added to detect the production of ornithine decarboxylase. Glucose is the fermentable carbohydrate, which during the initial stages of incubation, is fermented by the organisms with acid production, which results in colour change of the pH indicator (BCP) to yellow. The acidic condition also stimulates decarboxylase activity. If the organism produces the appropriate enzyme, i.e. decarboxylase, the amino acid (ornithine) in the medium is degraded, yielding a corresponding amine. Decarboxylation of ornithine yields putrescine. The production of this amine elevates the pH of the medium towards alkalinity, changing the color of the indicator from yellow to purple or violet. If the organism does not produce the appropriate enzyme, the medium remains acidic or yellow in colour. The broth was prepared according to manufacturer's specification by suspending 9.01 grams in 1000 ml distilled water. The medium was dissolved completely by heating on electric hot plate. About 5 ml of the broth was dispensed in test tubes and autoclaved at 121°C for 15 minutes. The tubes were allowed to cool in an upright position and then aseptically incubated with the test organisms. Each inoculated tube was overlaid with 2-3 ml of sterile (autoclaved) paraffin oil and incubated at 37°C for 24 hours. Ornithine positive isolates changed the broth to purple while the tubes containing ornithine negative isolates has no colour change (i.e remained yellow) [12].

2.5.10 Lysine decarboxylation test

Lysine decarboxylase broth was prepared according to manufacturer's specification by suspending 14.02 grams in 1000 ml distilled water. The medium was dissolved completely by heating on electric hot plate. About 5ml of the broth was dispensed in test tubes and autoclaved at 121°C for 15 minutes. The tubed broth was allowed to cool in an upright position and then aseptically incubated with the test organisms. Each inoculated tubed broth was overlaid with 2-3 ml of sterile (autoclaved) paraffin oil and incubated at 37°C for 2 to 4 days. Lysine positive isolates changed the broth to purple while the tubes containing ornithine negative isolates has no colour change (i.e remained yellow) [13].

2.5.11 Fermentation of sugars

The bacterial isolates were tested for the fermentation of sugars such as glucose, sucrose, mannitol and xylose. 1.0 g of each sugar was weighed and dispensed into different conical flasks and labeled. 2.5 g of peptone water was added into 100 ml of distilled water and 0.01 g of phenol red was added as the indicator. 5ml of each sugar was dispensed into 15 mls test tubes with Durham's tube introduced in upturned position into each test tube. Each test tube was corked and autoclaved for 15 minutes at 121°C. After autoclaving, the test tubes were allowed to cool after which bacterial isolates were aseptically inoculated into the sugar solution and incubated at 37°C for 2 to 5 days. Changes in the colour from red to yellow indicated production of acid, which implied utilization of the sugar by the bacterial isolate. Collection of gas bubbles in the Durham's tube indicated gas production [11].

2.6 Antibiotic Sensitivity Test

A 0.5 McFarland standard was prepared by mixing 0.05 mL of 1.175% barium chloride dihydrate ($\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$), with 9.95 mL of 1% sulfuric acid (H_2SO_4). 0.5McFarland corresponds to 1.5×10^8 , which is the bacterial turbidity required for antibiotic sensitivity test The 103 bacterial isolates isolated from human faecal storage cisterns were subjected to ten (10) commonly use commercially available antibiotics, viz: Augmentin (30 µg), Gentamycin (10 µg), Pefloxacin (10 µg), Ofloxacin (5 µg), Streptomycin (10 µg), Sulfomethoxazole-trimethoprim (30 µg), Chloramphenicol (30 µg); Sparfloxacin (10 µg); Ciprofloxacin (5 µg) and Amoxicillin (20 µg) [6].

Using Kirby-Bauer method, colonies from overnight culture of bacterial isolate, was aseptically picked and inoculated into test tube containing peptone water that had been autoclaved after it was prepared according to manufacturer's specification. The inoculated test tube was then incubated in a bacteriological incubator at 37°C for about 18 hours. Mueller-Hinton agar (Oxoid, UK) was prepared according to manufacturer's specification, autoclaved at 121°C for 15 minutes. The agar was allowed to cool to about 42°C and then poured into various petri dishes, reaching 4 mm. The agar was then allowed to congeal in the plates. The turbidity of the test tubes containing the incubated isolates (one isolate per tube) were compared to the prepared McFarland standard and diluted with normal saline when necessary. Sterile cotton bud was inserted into each tube and used to evenly swab the surface of the Mueller-Hinton agar contained in the petri dishes (three petri dishes to an isolate). This was done in less than 15 minutes after comparing with McFarland standard. After few minutes, antibiotic disks were then aseptically picked and placed on the petri dishes. The plates were then incubated at 37°C for 18 to 24 hours, after which the zone of inhibition for each isolates was then measured and recorded [14].

2.7 Statistical Analysis

Data obtained were subjected to two-way analysis of variance (ANOVA) and treatment means were separated using Duncan's New Multiple Range Test (DBMRT) at $P \leq 0.05$ level of significance with the aid of Statistical Package for Social Sciences (SPSS) version 23.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Mean colony forming unit of the bacterial isolates on MacConkey agar

The colony forming units of the bacterial Isolates from each location are shown (Fig. 1). Akure has the highest mean colony forming unit of Akure 42.44 ± 1.82^b , while Akungba has the least mean colony forming unit (35.00 ± 1.27^a).

3.1.2 Biochemical characterization of the isolates

All the 103 bacterial isolates were subjected to various Biochemical test. The results of the biochemical tests and corresponding

characterized isolates are shown (Table 1). The tests identified ten (10) bacterial organisms, viz: EC = *Escherichia coli*; EA= *Enterobacter aerogenes*; KP= *Klebsiella pneumoniae*; ST= *Salmonella typhi*; PA = *Pseudomonas aeruginosa*; SD = *Shigella dysenteriae*; PM= *Proteus mirabilis*; CK=*Citrobacter koseri*; PF= *Providentia alcalifaciens*; and KO = *Klebsiella oxytoca*. Table 2 shows the distribution of the isolates with respect to location. The table shows that *E. coli*, (32.04%), occurred more than any organisms identified.

3.1.3 Antibiotic sensitivity in the bacterial isolates

By following standard methods, the isolates were subject to ten (10) antibiotics, using impregnated antibiotic disks. Fig. 2 shows the zone of inhibition of the isolates with respect to locations. Table 3 shows the mean zone of inhibition of the isolates to the antibiotics used. The list zone of inhibition (0.78 ± 0.68^a) was found in *C. koseri* against Augmentin; While the highest zone of inhibition (20.83 ± 0.01^b) was found in *P. aeruginosa* against Ciprofloxacin. Table 4 shows the antibiotic resistance patterns of the bacterial isolates: the isolates were least sensitive to Augmentin, where 98 (95.15%) of the isolates showed resistance; the highest sensitive was found against Ciprofloxacin, where 28 (27.18%) of the isolates showed resistance. Table 5 shows degree of resistance based on the classes of antibiotics to which the isolates showed resistance. From the table, it can be seen that only 2 (1.94%) isolates showed resistance to two classes of antibiotics whereas as many 38 (36.89%) isolates showed resistance to the whole five classes of antibiotics used.

3.2 Discussion

The colony forming units (on MaConckey agar) of bacterial isolates from the collected samples were found to be very similar in the samples collected from Akungba (35.00 ± 1.27^a cfu/g) and Ore (35.25 ± 1.49^a cfu/g); and Owo (36.25 ± 1.51^a cfu/g) and Ondo (36.43 ± 1.38^a cfu/g). In fact, the colony forming units of the bacterial isolates from these four locations could be regarded as similar. Only the colony forming unit of the bacterial isolates from Akure samples (42.44 ± 1.82^b cfu/g) was totally different with at least a difference of 7 when compared distinctly with other locations'. These similarities and differences could be as a result the variation in sample sizes from the locations involved.

Of the 103 bacterial isolates in this study, 26 (25.2%) were isolated from faecal samples from Akure; same number of isolates, 21 (20.4%) were isolated from Akungba and Ore; 18 (17.5%) from Owo; and 17 (16.5%) from Ondo. Biochemical characterization showed that the bacterial isolates were *Escherichia coli* (32.04%), *Enterobacter aerogenes* (14.56%), *Klebsiella pneumoniae* (12.62%), *Salmonella typhi* (11.65%), *Pseudomonas aeruginosa* (7.77%), *Shigella dysenteriae* (6.80%), *Proteus mirabilis* (5.83%), *Citrobacter koseri* (3.88%), *Providentia alcalifaciens* (2.91%) and *Klebsiella oxytoca* (1.94%). The high number of *E. coli* is likely due to the fact that the bacterium is an aerobic intestinal organism that are passed out with fresh faeces; and it is in conformity with the work of Islam, et al. [9], who isolated *E. coli* as the most occurring bacteria from human faecal samples. The presence of *Pseudomonas aeruginosa*, which is not an intestinal bacterium, in relatively large numbers can be linked to the versatility of the bacterium to adapt in various environment. The 7.77% *Pseudomonas aeruginosa* population in this work tallies with the one isolated in the work of Onuoha [15]. *Escherichia coli*, *Citrobacter koseri*, *Salmonella typhi* and *Enterobacter aerogenes* (synonym: *Klebsiella aerogenes*), identified in this study, have earlier been identified by Nworie, et al. [16], who isolated same bacteria from public toilets in Nigeria. ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp.) pathogens are responsible for increase in antimicrobial-resistant infections worldwide and have been rated important pathogens by WHO [17]. Therefore, the isolation of *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Enterobacter aerogenes* in this research means that the toilet is an environment that can enhance the propagation of important pathogens that are inimical to human health. The work of Kaur, et al. [18], who isolated pathogenic bacteria from toilet tap waters, further lend credence to this. *Proteus mirabilis* and *Escherichia coli* are major causes of urinary tract infection; *Pseudomonas aeruginosa* is known to form biofilm in indwelling medical devices like catheter; *Klebsiella pneumoniae* and *K. oxytoca* have been isolated from upper respiratory tract infection patients [19]; *Citrobacter koseri* has been identified as a microbial cause of meningitis and cerebral abscess [20]; while *Providentia alcalifaciens* is associated with traveller's diarrhoea [21], all of

Table 1. Biochemical characterization of the bacterial isolates

Probable Bacteria	Triple sugar iron			GR	MS	C _o MAC	MR	VP	CAT	OXI	COA	IND	URS	MOT	CIT	ORNITINE	LYSINE	SUCROSE	MANNITOL	GLUCOSE	XYLOSE
	Slant	Butt	H ₂ S																		
EC	ACID	ACID	-	-	R	PK	+	-	+	-	-	+	-	MT	-	+	+	+	+	+	+
EA	ACID	ACID	-	-	R	PK	-	+	+	-	-	-	-	MT	+	+	+	+	-	+	+
KP	ACID	ACID	-	-	R	PK	-	+	+	-	-	-	+	NM	+	-	+	+	+	+	+
PA	ALK	ALK	-	-	R	CL	-	-	+	+	-	-	-	MT	+	-	-	-	+	+	-
ST	ACID	ALK	+	-	R	CL	+	-	+	-	-	-	-	MT	-	-	+	-	+	+	+
SD	ALK	ACID	-	-	R	CL	+	-	+	-	-	+	-	NM	-	-	-	-	-	+	-
PM	ACID	ALK	+	-	R	CL	+	-	+	-	-	-	+	MT	+	+	-	-	-	+	+
CK	ACID	ACID	-	-	R	PK	+	-	+	-	-	+	+	MT	+	-	-	+	+	+	+
KO	ACID	ACID	-	-	R	PK	-	+	+	-	-	+	+	NM	+	-	+	+	+	+	-
PF	ALK	ACID	-	-	R	CL	+	-	+	-	-	+	-	MT	+	-	-	-	-	+	+

EC= *Escherichia coli*; EA= *Enterobacter aerogenes*; KP= *Klebsiella pneumoniae*; ST= *Salmonella typhi*; PA= *Pseudomonas aeruginosa*; SD= *Shigella dysenteriae*; PM= *Proteus mirabilis*; CK= *Citrobacter koseri*; PF= *Providentia alcalifaciens*; KO= *Klebsiella oxytoca*, GR=Gram reaction, MS=Microscopic shape, CoMAC=Color on MacConkey agar; MR=Methyl Red; VP=Voges praskauer; CAT=Catalase; OXI=Oxidase; COA=Coagulase; IND=Indole; URS=urease; MOT=Motility, CIT=Citrate; PK=Pink; CL=Colourless; ALK=Alkaline; MT=Motility; NM=Non-motile

which were isolated from faecal samples used in this study. It is important to state that there is a continuing high burden of typhoid fever in many

parts of the world and a rapid increase in the emergence and spread of antimicrobial resistant strains of *S. typhi* [22].




Fig. 1. Mean colony forming unit per gram of sample on MaConckey agar, with respect to location




Fig. 2. Mean zone of inhibition (with error bars) of the antibiotics disks to the isolates from each location. Each value is a mean of 3 replicates

AU=Augmentin; GN=Gentamycin; PEF=Pefloxacin; OFX=Ofloxacin; S=Septin; SXT=Sulphomethazole-trimethoprim; CH=Chloramphenicol; SP=Sparfloxacin; CPX= Ciprofloxacin; AM=Ampicillin

Table 2. Percentage distribution pattern of the bacterial isolates with respect to locations

Isolate	AKUNGBA	OWO	AKURE	ONDO	ORE	N/O	% DIS
<i>E. coli</i>	6	5	10	5	7	33	32.04
<i>E. aerogenes</i>	5	2	3	3	2	15	14.56
<i>K. pneumoniae</i>	1	2	4	3	3	13	12.62
<i>S. typhi</i>	2	2	3	2	3	12	11.65
<i>P. aeruginosa</i>	2	1	2	2	1	8	7.77
<i>S. dysenteriae</i>	2	1	2	0	2	7	6.80
<i>P. mirabilis</i>	0	2	1	2	1	6	5.83
<i>C. koseri</i>	1	1	0	0	2	4	3.88
<i>P. alcalifaciens</i>	1	2	0	0	0	3	2.91
<i>K. oxytoca</i>	1	0	1	0	0	2	1.94
Total	21(20.4%)	18(17.5%)	26(25.2%)	17(16.5%)	21(20.4%)	103	100

EC= *Escherichia coli*; EA= *Enterobacter aerogenes*; KP= *Klebsiella pneumoniae*; ST= *Salmonella typhi*; PA= *Pseudomonas aeruginosa*; SD= *Shigella dysenteriae*; PM= *Proteus mirabilis*; CK= *Citrobacter koseri*; PF= *Providentia alcalifaciens*; KO= *Klebsiella oxytoca*. N/O=Number of occurrence; DIS= Distribution

There were variations in the antibiotic sensitivity profile of the bacterial isolates. All the bacteria in this study showed resistance to multiple antibiotics. For example, *Escherichia coli* has a minimum zone of inhibition of 3.27 ± 0.02^a , which was showed against Augmentin and a maximum zone of inhibition of 18.90 ± 0.09^b , showed against Ciprofloxacin. This means that following [6] zone of inhibition interpretative criteria, none of the *E. coli* in this study was sensitive to any of the 10 antibiotics used. In fact, 90.91% of the *E. coli* tested showed resistance to Augmentin. The resistance of the isolates to the various antibiotics used varies from 27.18% to Ciprofloxacin, 33.98% to Ofloxacin, 47.57% to Sulphomethazole-Trimethoprim, 56.31% to Streptomycin, 66.99% to Chloramphenicol, 74.76% to Pefloxacin, 75.73% to Sparfloxacin, 75.73% to Gentamycin, 85.44% to Ampicillin and 95.15 to Augmentin. The least resistance, 27.18%, was found against Ciprofloxacin while the highest, 95.15, was found against Augmentin. This is a little more above what was observed in the findings of Sohn, et al. [23] - an indication that antibiotic resistant bacteria are on the increase. The relatively low number of Isolates which showed resistance to ciprofloxacin may be as a result of the drug's pronounced potency against Gram negative enterobacteriaceae, by interfering with their DNA gyrase [24] and thereby hindering a complete replication. Meanwhile, [25] findings also showed that enterobacteriaceae are relative sensitive to ciprofloxacin. More worrisome is the high level of multidrug resistance isolates. At least 84.46% of the isolates are multidrug resistant. That is, they showed resistance to three or more classes of antibiotics [26]. Of these multidrug resistant isolates, 43.68% showed resistance to all the five groups of antibiotics used. This is very disturbing

considering the fact that O'Neill [4] has predicted a post antibiotic era against year 2050 where 10 million people are expected to die yearly due to antibiotic resistant bacteria.

The high distribution, as shown by this study, of multidrug resistant bacteria in faecal storage cisterns, is a big concern. This can be linked to many factors. For example, discarding unused drugs into toilets can contribute to increase in drug resistance among bacteria. Some of the toilets used in this study are two-in-one, that is people can also bath in the toilet. Meanwhile the waste water after bathing goes to the same tank with the faeces. Since there are many antibacterial soaps used for bathing, it is not unlikely that they will contribute to antibacterial resistance among the toilet bacteria. Bacteria in the toilets can find their way, via vertical or horizontal flow, particularly during raining seasons, into water aquifers. Waters from this aquifers will enhance the spread of pathogenic and antibiotic resistance bacteria if used without appropriate sterilization.

In the same way, cockroaches can also propagate the spread of multidrug resistance and ESBL-producing bacteria from the toilet. Cockroaches are insects with long antennae and legs, feeding by scavenging. They are one of the most significant and objectionable pests found in apartments, homes, food-handling establishments, hospitals and health care facilities worldwide. Indoor species, especially the German cockroach, exploit conditions associated with high-density human populations and impoverished living conditions [27,28]. Similar bacteria isolated from human faecal samples in this study have also been reportedly isolated from various parts of of cockroaches by

Table 3. Zones of inhibitions of the isolated bacteria to the antibiotics used

Tested organism	AU	GN	PEF	OFX	S	SXT	CH	SP	CPX	AM
<i>Escherichia coli</i>	3.27±0.02 ^a	9.76±0.03 ^c	13.87±0.01 ^h	15.06±0.03 ⁱ	10.88±0.01 ^f	10.06±0.06 ^d	10.52±0.04 ^e	11.03±0.03 ^g	18.90±0.09 ^b	9.38±0.08 ^j
<i>Enterobacter aerogenes</i>	3.16±1.16 ^a	8.25±0.01 ^c	10.03±0.03 ^d	12.69±0.01 ^e	7.28±0.01 ^b	7.07±0.01 ^b	8.49±0.01 ^c	9.49±0.00 ^d	17.89±0.10 ^b	6.80±0.00 ^f
<i>Klebsiella pneumoniae</i>	3.81±0.01 ^a	7.93±0.03 ^b	11.34±0.30 ^g	14.52±0.01 ^h	9.81±0.16 ^e	9.07±0.06 ^d	10.65±0.01 ^f	10.70±0.02 ^f	19.15±0.01 ^c	8.13±0.02 ^j
<i>Salmonella typhi</i>	1.37±0.01 ^a	7.09±0.02 ^c	10.26±0.01 ^g	12.09±0.01 ⁱ	8.91±0.08 ^e	8.11±0.00 ^d	10.17±0.01 ^f	10.74±0.04 ^h	17.27±0.03 ^b	6.93±0.06 ^j
<i>Pseudomonas aeruginosa</i>	7.50±0.03 ^a	11.45±0.05 ^e	15.71±0.03 ^h	16.79±0.00 ⁱ	11.39±0.02 ^d	12.45±0.05 ^g	11.26±0.02 ^c	11.51±0.01 ^f	20.83±0.01 ^b	9.59±0.01 ^j
<i>Shigella dysenteriae</i>	2.52±0.01 ^a	9.06±0.01 ^c	11.85±0.01 ^g	15.86±0.12 ^h	11.39±0.01 ^e	11.91±0.00 ^g	11.81±0.01 ^f	11.05±0.05 ^d	16.25±0.01 ^b	8.68±0.02 ^j
<i>Proteus mirabilis</i>	2.56±0.00 ^a	9.50±0.01 ^d	11.94±0.01 ^f	16.94±0.00 ^h	9.11±0.00 ^c	13.08±0.02 ^g	11.00±0.00 ^e	11.00±0.02 ^e	17.29±0.01 ^b	7.78±0.00 ^j
<i>Citrobacter koseri</i>	0.78±0.68 ^a	7.94±0.04 ^b	11.42±0.01 ^e	13.34±0.01 ^h	12.33±0.02 ^f	12.83±0.02 ^g	9.28±0.02 ^d	15.86±0.02 ⁱ	18.34±0.01 ^c	8.50±0.00 ^j
<i>Providentia alcalifaciens</i>	5.18±0.07 ^b	9.93±0.13 ^g	9.33±0.01 ^e	12.34±0.01 ^h	9.80±0.07 ^f	6.91±0.08 ^c	9.05±0.09 ^d	14.34±0.02 ⁱ	20.32±0.03 ^a	4.03±0.06 ^j
<i>Klebsiella oxytoca</i>	0.72±0.62 ^a	4.13±0.15 ^c	5.00±0.00 ^d	8.00±0.00 ^f	2.19±0.02 ^b	2.03±0.06 ^b	7.82±0.01 ^f	6.68±0.02 ^e	13.07±0.06 ^a	0.75±0.65 ^g

Each value is a mean of three replicates. Values in the same column with same superscripts are not significantly different @ P ≤ 0.05. Key: AU=Augmentin; GN=Gentamycin; PEF=Pefloxacin; OFX=Ofloxacin; S=Seprin; SXT=Sulphomethazole-trimethoprim; CH=Chloramphenicol; SP=Sparfloxacin; CPX= Ciprofloxacin; AM=Ampicillin

Table 4. Antibiotic resistance pattern of the bacterial isolates [n (%)] to the various antibiotics used

Bacterial isolates	I.T	AU	GN	PEF	OFX	S	SXT	CH	SP	CPX	AM
<i>E. coli</i>	33	30 (90.91)	21(63.64)	22(66.67)	9(27.27)	15(45.45)	14(42.42)	22(66.67)	25(75.76)	9(27.27)	27(81.82)
<i>E. aerogenes</i>	15	15(100)	13(86.67)	12(80.00)	7(46.67)	10(66.67)	10(66.67)	9(60.00)	13(86.67)	5(33.33)	11(73.33)
<i>K. pneumoniae</i>	13	12(92.31)	10(76.92)	11(84.64)	3(23.08)	8(61.54)	7(53.85)	8(61.54)	12(92.31)	2(15.38)	13(100)
<i>S. typhi</i>	12	12(100)	9(75.00)	10(83.33)	7(53.33)	7(53.33)	9(75.00)	10(83.33)	4(33.33)	4(33.33)	11(91.67)
<i>P. aeruginosa</i>	8	7(87.50)	6(75.00)	5(62.50)	1(12.50)	4(50.00)	3(37.50)	5(62.50)	8(100)	2(25.00)	6(75.00)
<i>S. dysenteriae</i>	7	7(100)	7(100)	4(57.14)	2(28.57)	3(42.86)	1(14.29)	5(71.43)	6(85.71)	2(28.57)	5(71.43)
<i>P. mirabilis</i>	6	6(100)	4(66.67)	5(83.33)	0(0.00)	5(83.33)	1(16.67)	3(50.00)	4(66.67)	2(33.33)	5(83.33)
<i>C. koseri</i>	4	4(100)	3(75.00)	3(75.00)	3(75.00)	3(75.00)	0(0.00)	3(75.00)	2(50.00)	0(0.00)	4(100)
<i>P. alcalifaciens</i>	3	3(100)	3(100)	3(100)	2(66.67)	1(33.33)	2(66.67)	3(100)	2(66.67)	0(0.00)	3(100)
<i>K. oxytoca</i>	2	2(100)	2(100)	2(100)	1(50.00)	2(100)	2(100)	1(50.00)	2(100)	2(100)	2(100)
Total	103	98(95.15)	78(75.73)	77(74.76)	35(33.98)	58(56.31)	49(47.57)	69(66.99)	78(75.73)	28(27.18)	88(85.44)

I.T= Isolates tested; AU=Augmentin; GN=Gentamycin; PEF=Pefloxacin; OFX=Ofloxacin; S=Seprin; SXT=Sulphomethazole-trimethoprim; CH=Chloramphenicol; SP=Sparfloxacin; CPX= Ciprofloxacin; AM=Ampicillin

many authors in Nigeria [29,30,31] and Ethiopia [28]. The crevices in underground faecal storage cisterns make it more easier for cockroaches to spread antibiotic resistant bacteria in communities and hospital environments. Since cockroaches can also inhabit the dark areas in water wells, they may as well spread drug resistance bacteria from the toilets to the well water. Meanwhile multidrug resistant bacteria have been isolated from wells and underground water storage tanks in Nigeria [32,33]; from surface waters [34]; as well as from toilet indoor airs and toilet door handles [35,36].

Table 5. Number of antibiotics group and isolates that showed resistance to (a) given group(s)

No of antibiotics group	No of resistant isolates (%)
One	2 (1.94)
Two	14 (13.59)
Three	16 (15.33)
Four	33 (32.02)
Five	38 (36.89)

The spread of multidrug resistant bacteria from faecal cisterns crevices within community and hospital environment is not limited to cockroach vectors. Flies, rats and mouse are potential household vectors that can spread pathogenic and antibiotic resistant bacteria within communities. House flies (*Musca domestica*) have been known as a mechanical vector in spreading infectious diseases such as cholera, shigellosis, salmonellosis and skin infections. House flies are able to transport pathogenic agents by attaching them to their mouth, body surface, foot and wings [37]. Usui, et al. [38] reported flies carrying beta-lactamase genes. In another research by Petridis, et al. [39], shiga toxin producing genes was found in *E. coli* isolated from houseflies. In yet another research carried out differently by Liu, et al. [40], Usui, et al. [38], Fukuda, et al. [41 and Zhang, et al. [42], multidrug resistant Gram negative enteric bacteria; plasmids carrying antibiotic resistant genes; Cephalosporins resistance genes; and colistin resistance genes respectively were found in bacteria isolated from houseflies.

4. CONCLUSION

Findings from this study revealed that multiple-antibiotic resistant bacteria are in high distribution in faecal storage cisterns in Ondo State. And they can be transmitted to human by formite, animal vectors and water. Consequently,

adequate management of faecal storage cisterns is important. Further research is however needed to relate the bacteria isolated from the storage cisterns with bacteria present in water sources within the perimeters of the cisterns.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Laxminarayan R, Duse A, Wattal C, Zaidi AK, Wertheim HF, Sumpradit N, Vlieghe E, Hara GL, Gould IM, Goossens H, Greko C. Antibiotic resistance—the need for global solutions. *The Lancet Infectious Diseases*. 2013;13(12):1057-98.
2. Laxminarayan R, Heymann DL. Challenges of drug resistance in the developing world. *BMJ*. 2012;344:e1567.
3. World Health Organization. Antimicrobial resistance and primary health care: Brief. World Health Organization; 2018.
4. O'Neill J. The Wellcome Trust and HM Government, Securing New Drugs for Future Generations, The Pipeline of Antibiotics; 2015.
5. Nigeria Centre for Disease Control. Antimicrobial use and resistance in Nigeria. Situation Analysis and Recommendations. Produced by Federal Ministries of Agriculture, Environment and Health. Coordinated by Nigeria Centre for Disease Control; 2017.
6. Clinical and Laboratory Standard Institute (CLSI 2018 M100-28). Performance Standards for Antimicrobial Susceptibility Testing; 2018.
7. Gerhardt A, Hammer TR, Balluff C, Mucha H, Hofer D. A model of the transmission of micro-organisms in a public setting and its correlation to pathogen infection risks. *Journal of Applied Microbiology*. 2012;112(3):614-21.
8. Brown A, Smith H. Benson's microbiological applications, laboratory manual in general microbiology, Short Version. McGraw-Hill Education; 2014.
9. Islam MA, Amin MB, Roy S, Asaduzzaman M, Islam MR, Navab-Daneshmand T, Mattioli MC, Kile ML, Levy K, Julian TR. Fecal colonization with multidrug-resistant *E. coli* among healthy infants in rural Bangladesh. *Frontiers in Microbiology*. 2019;10:640.

10. Cheesbrough M. District Laboratory Practice in Tropical Countries. Cambridge University Press; 2006.
11. Fawole MO, Oso BA. Characterization of bacteria: Laboratory manual of microbiology. Spectrum Book Ltd., Ibadan, Nigeria. 2004;24-33.
12. McFaddin JF. Biochemical tests for the identification of medical bacteria. 2000; 170–182.
13. Jorgensen JH, Pfaller MA, Carroll KC, Funke G, Landry ML, Richter SS, Warnock, DW. Manual of clinical microbiology, 11th Edition. 2015;1.
14. Howe RA, Andrews JM. BSAC standardized disc susceptibility testing method (version 11). Journal of Antimicrobial Chemotherapy. 2012;67(12): 2783-4.
15. Onuoha SC. The prevalence of antibiotic resistant diarrhoegenic bacterial species in surface waters, South Eastern Nigeria. Ethiopian Journal of Health Sciences. 2017;27(4):319-30.
16. Nworie A, Ayeni JA, Eze UA, Azi SO. Bacterial contamination of door handles/knobs in selected public conveniences in Abuja metropolis, Nigeria: A public health threat. Continental Journal of Medical Research. 2012;6(1):7-11.
17. Navidinia M. The clinical importance of emerging ESKAPE pathogens in nosocomial infections. Archives of Advanced in Bioscience. 2016;7(3):43-57.
18. Kaur R, Singh D, Kesavan AK, Kaur R. Molecular characterization and antimicrobial susceptibility of bacterial isolates present in tap water of public toilets. International Health; 2019.
19. Motayo BO, Ogiogwa JI, Aboderin BW, Okerentugba PO, Innocent-Adiele HC, Nwanze JC, Onoh CC, Okonko IO. Bacteriological review of multi-drug resistance (MDR) pathogens involved in respiratory tract infections (RTIs) in Abeokuta, Nigeria. Researcher. 2012;4(5):49-55.
20. Marecos CV, Ferreira M, Ferreira MM, Barroso MR. Sepsis, meningitis and cerebral abscesses caused by *Citrobacter koseri*. Case Reports. 2012; bcr1020114941.
21. Shima A, Hinenoya A, Asakura M, Sugimoto N, Tsukamoto T, Ito H, Nagita A, Faruque SM, Yamasaki S. Molecular characterizations of cytolethal distending toxin produced by *Providencia alcalifaciens* strains isolated from patients with diarrhea. Infection and Immunity. 2012;80(4):1323-32.
22. World Health Organization. Typhoid vaccines: WHO position paper, March 2018–Recommendations. Vaccine. 2019; 37(2):214-6.
23. Sohn KM, Kang CI, Joo EJ, Ha YE, Chung DR, Peck KR, Lee NY, Song JH. Epidemiology of ciprofloxacin resistance and its relationship to extended-spectrum β -lactamase production in *Proteus mirabilis* bacteremia. The Korean Journal of Internal Medicine. 2011;26(1):89.
24. Wang F, Raval Y, Tzeng TR, Anker JN. X-Ray excited luminescence chemical imaging of bacterial growth on surfaces implanted in tissue. Advanced Healthcare Materials. 2015;4(6):903-10.
25. Temitope OO, Emmanuel KT. Antibiotic Resistant Pattern of Bacteria Isolated from Faecal Pellets of *Supella longipalpa* (Cockroach) Found in Akungba Akoko, Ondo State, Nigeria. Journal of Analytical Techniques and Research. 2019;1:010-24.
26. Falagas ME, Karageorgopoulos DE. Pandrug resistance (PDR), extensive drug resistance (XDR) and multidrug resistance (MDR) among Gram-negative bacilli: Need for international harmonization in terminology. Clinical Infectious Diseases. 2008;46(7):1121-2.
27. Bonnefoy X, Kampen H, Sweeney K. Public health significance of urban pests. World Health Organization; 2008.
28. Moges F, Eshetie S, Endris M, Huruy K, Muluye D, Feleke T, Ayalew G, Nagappan R. Cockroaches as a source of high bacterial pathogens with multidrug resistant strains in Gondar town, Ethiopia. BioMed Research International. 2016; 2016.
29. Bala AY, Sule H. Vectorial potential of cockroaches in transmitting parasites of medical importance in Arkilla, Sokoto, Nigeria. Nigerian Journal of Basic and Applied Sciences. 2012;20(2):111-5.
30. Iboh CI, Etim LB, Abraham JT, Ajang RO. Bacteria and parasites infestation of cockroaches in a developing community, South Eastern, Nigeria. Int. J. Bacteriol. Res. 2014;2(5):45-8.
31. Isaac C, Orue PO, Iyamu MI, Ehiaghe JI, Isaac O. Comparative analysis of pathogenic organisms in cockroaches from different community settings in Edo State,

- Nigeria. The Korean Journal of Parasitology. 2014;52(2):177.
32. Ajayi AO, Adejumo TO. Microbiological assessment and some physico-chemical properties of water sources in Akungba-Akoko, Nigeria. Journal of Toxicology and Environmental Health Sciences. 2011; 3(13):342-6.
 33. Oluyeye JO, Dada AC, Odeyemi AT. Incidence of multiple antibiotic resistant Gram-negative bacteria isolated from surface and underground water sources in south western region of Nigeria. Water Science and Technology. 2009;59(10): 1929-36.
 34. Ayandele AA, Ajala OO, Oyekemi SA, Awotunde MO. Microbiological evaluation and antimicrobial resistant pattern of bacteria isolated from surface drinking water sources in Ogbomoso, Oyo State, Nigeria. Nigerian Journal of Biotechnology. 2019;36(1):17-26.
 35. Alonge OO, Auwal BM, Aboh MI. Bacterial contamination of toilet door handles on Baze University campus Abuja Nigeria. African Journal of Clinical and Experimental Microbiology. 2019;20(1):35-41.
 36. Ohagim PI, Ikon GM, Matthew PC, Ohagim GA. Microbiological assessment of indoor air in public toilets across selected motor parks in Owerri Metropolis, Nigeria. J Microbiol Exp. 2017;5(6):00166.
 37. Nazari M, Mehrabi T, Hosseini SM, Alikhani MY. Bacterial contamination of adult house flies (*Musca domestica*) and sensitivity of these bacteria to various antibiotics, captured from Hamadan City, Iran. Journal of Clinical and Diagnostic Research: JCDR. 2017;11(4):DC04.
 38. Usui M, Shirakawa T, Fukuda A, Tamura Y. The role of flies in disseminating plasmids with antimicrobial-resistance genes between farms. Microbial Drug Resistance. 2015;21(5):562-9.
 39. Petridis M, Bagdasarian M, Waldor MK, Walker E. Horizontal transfer of Shiga toxin and antibiotic resistance genes among *Escherichia coli* strains in house fly (Diptera: Muscidae) gut. Journal of Medical Entomology. 2014;43(2):288-95.
 40. Liu Y, Yang Y, Zhao F, Fan X, Zhong W, Qiao D, Cao Y. Multi-drug resistant gram-negative enteric bacteria isolated from flies at Chengdu Airport, China. Southeast Asian J Trop Med Public Health. 2013; 44(6):988-6.
 41. Fukuda A, Usui M, Okubo T, Tamura Y. Horizontal transfer of plasmid-mediated cephalosporin resistance genes in the intestine of houseflies (*Musca domestica*). Microbial Drug Resistance. 2016;22(4): 336-41.
 42. Zhang J, Wang J, Chen L, Yassin AK, Kelly P, Butaye P, Li J, Gong J, Cattley R, Qi K, Wang C. Housefly (*Musca domestica*) and blow fly (*Protophormia terraenovae*) as vectors of bacteria carrying colistin resistance genes. Appl. Environ. Microbiol. 2018;84(1):e01736-17.

© 2020 Onifade and Abi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/54412>